

# Isolation of bis(copper) key intermediates in Cu-catalyzed azide-alkyne “click reaction”

Liqun Jin, Daniel R. Tolentino, Mohand Melaimi, Guy Bertrand\*

The copper-catalyzed 1,3-dipolar cycloaddition of an azide to a terminal alkyne (CuAAC) is one of the most popular chemical transformations, with applications ranging from material to life sciences. However, despite many mechanistic studies, direct observation of key components of the catalytic cycle is still missing. Initially, mononuclear species were thought to be the active catalysts, but later on, dinuclear complexes came to the front. We report the isolation of both a previously postulated  $\pi,\sigma$ -bis(copper) acetylide and a hitherto never-mentioned bis(metallated) triazole complex. We also demonstrate that although mono- and bis-copper complexes promote the CuAAC reaction, the dinuclear species are involved in the kinetically favored pathway.

The copper-catalyzed azide-alkyne cycloaddition (CuAAC) (1, 2) is the flagship of the “click chemistry” arsenal (3, 4). Because of the straightforward procedures, the high functional group tolerance, and the usually high yields, CuAAC has affected all fields of chemistry from material (5–8) to life sciences (9–13); however, the mechanism of the CuAAC is still under debate (14, 15). The first postulated mechanism involved mono-copper complexes, although more recently, the participation of bis(copper) complexes has been suggested on the basis of kinetic experiments (16) and computational investigations (17–19). In 2013, Fokin and colleagues published a very elegant mechanistic study and stated that “monomeric copper-acetylide complexes  $\mathbf{1}_{Cu}$  are not reactive towards organic azides unless an exogenous copper catalyst is added” (20). The marked difference with the previously accepted mechanism is the introduction of the cationic  $\pi,\sigma$ -bis(copper) acetylide complexes of type  $\mathbf{1}_{Cu2}$  (Fig. 1) as the “catalytically active complex,” instead of  $\mathbf{1}_{Cu}$ . However, such a dinuclear species was qualified as a “non-isolable and highly reactive intermediate.” We have recently shown that the strong  $\sigma$ -donating and  $\pi$ -accepting properties of cyclic (alkyl)(amino) carbenes (CAACs) (21–23) allow for the isolation of electron-deficient (24), electron-rich (25), and even paramagnetic species (26). This prompted us to attempt using CAAC ligands to stabilize elusive species involved in the CuAAC catalytic cycle. Here, we report the isolation of a  $\pi,\sigma$ -bis(copper) complex of type  $\mathbf{1}_{Cu2}$  and of an unprecedented catalytically active 3,5-bis(metallated) triazole  $\mathbf{2}_{Cu2}$ . Thanks to the unusual stability of these complexes, we compare the mono- and dinuclear-based mechanisms and bring unambiguous evidence that the latter is indeed kinetically favored.

We first prepared the monomeric (27, 28) copper complex  $\mathbf{1}_{Cu}$  (figs. S7 and S8) by reacting lithium phenylacetylide with (CAAC)CuOAc (figs. S1 and S2) in tetrahydrofuran (THF) (Fig. 1A). We found that  $\mathbf{1}_{Cu}$  cleanly reacted with (CAAC)CuOTf (figs. S3 to S6) in methylene chloride, affording the desired cationic dinuclear complex  $\mathbf{1}_{Cu2}$  (figs. S9 and S10), which was isolated as a white solid in 95% yield. The  $^1\text{H}$  and  $^{13}\text{C}$  nuclear magnetic resonance (NMR) spectra show only one set of signals for both CAAC ligands, even at temperatures as low as  $-80^\circ\text{C}$ , which is indicative of a very fast exchange between the two “Cu(CAAC)” units. However, in the solid state, a single crystal x-ray diffraction study shows two different types of coordination,  $\pi$  and  $\sigma$  (Fig. 1C, left). The Cu-Cu distance [2.9387(4) Å] is slightly longer than

those computationally predicted by Fokin and colleagues for model compounds (2.85 to 2.88 Å) (19).  $\mathbf{1}_{Cu2}$  proved to be air-stable and thermally robust [melting point (mp):  $174^\circ\text{C}$ ].

Despite its thermodynamic stability,  $\mathbf{1}_{Cu2}$  reacts cleanly with benzyl azide in methylene chloride, affording  $\mathbf{2}_{Cu2}$  (figs. S11 and S12), which was isolated in 87% yield as an air- and moisture-stable, pale yellow solid.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra show the presence of one benzyl and one phenyl group, as well as two magnetically nonequivalent CAAC moieties. X-ray diffraction studies of the corresponding tetrafluoroborate salt confirmed the 3,5-dimetallated-1,2,3-triazole structure  $\mathbf{2}_{Cu2}$  (Fig. 1C, center).

Because dinuclear complexes of type  $\mathbf{2}_{Cu2}$  had never been postulated in any CuAAC mechanism, the next question was whether it was involved in the catalytic cycle. Addition of a stoichiometric amount of phenyl acetylene to  $\mathbf{2}_{Cu2}$  led, after 5 hours, to the quantitative release of the 1-benzyl-4-phenyl-1,2,3-triazole  $\mathbf{3}$  accompanied by the regeneration of the bis-copper complex  $\mathbf{1}_{Cu2}$ . As an additional confirmation, the same reaction was repeated using  $\mathbf{2}_{Cu2}$  and a different alkyne ( $\text{Ph}_2\text{CHC}\equiv\text{CH}$ ), which resulted in the isolation of  $\mathbf{3}$  along with the corresponding  $\pi,\sigma$ -bis-copper complex  $\mathbf{1}_{Cu2}$  (Fig. 1C, right; figs. S13 and S14) in quantitative yields.

After having shown that the bis-copper complex  $\mathbf{1}_{Cu2}$  could be part of the catalytic cycle, we turned our attention to the corresponding mono-copper complex  $\mathbf{1}_{Cu}$ . We found that a stoichiometric amount of benzyl azide slowly reacted with complex  $\mathbf{1}_{Cu}$ , leading, after 16 hours, to the C-metallated heterocycle  $\mathbf{2}_{Cu}$  (figs. S15 and S16), which was isolated as a pale yellow solid in 61% yield (29). Addition of one equivalent of phenyl acetylene to  $\mathbf{2}_{Cu}$  led, after 5 hours, to the quantitative isolation of triazole  $\mathbf{3}$  and regeneration of the mono-copper complex  $\mathbf{1}_{Cu}$ .


These results as a whole suggest that both the mono- and bis-copper complexes could lead to the triazole  $\mathbf{3}$ , at least under stoichiometric conditions. However, the kinetic profile of the reactions of  $\mathbf{1}_{Cu}$  and  $\mathbf{1}_{Cu2}$  with benzyl azide revealed critical rate acceleration when the bimetallic complex is used [ $k_{\text{obs}}(\mathbf{1}_{Cu2})/k_{\text{obs}}(\mathbf{1}_{Cu}) > 94$ ] (Fig. 1B and fig. S17) (30). In contrast, the protodemetalation of  $\mathbf{2}_{Cu2}$  and  $\mathbf{2}_{Cu}$  affording triazole  $\mathbf{3}$  and regenerating  $\mathbf{1}_{Cu2}$  and  $\mathbf{1}_{Cu}$ , respectively, proceeds at a similar rate (fig. S18). It should be noted that in both cases, the protodemetalation arises from the reaction with the terminal alkyne and, thus, does not necessitate an external Brønsted acid. These reactions lead back to the acetylide complexes  $\mathbf{1}_{Cu2}$  and  $\mathbf{1}_{Cu}$ , which exclude the copper precatalyst (CAAC)CuOTf from the catalytic cycle.

The experiments discussed above reproduce parts of both postulated catalytic cycles under stoichiometric conditions. Each step proved to


UCSD-CNRS Joint Research Chemistry Laboratory (UMI 3555), Department of Chemistry, University of California San Diego, La Jolla, CA 92093–0343, USA.

\*Corresponding author. E-mail: guybertrand@ucsd.edu





**Fig. 2. Kinetic profiles of the CuAAC reaction of phenyl acetylene with benzyl azide using CAAC-supported mono- and bis(copper) catalysts.** (A) The kinetic profiles of the catalytic reaction of phenyl acetylene with benzyl azide demonstrate the superior catalytic activity of the dinuclear complexes  $1_{Cu_2a}$  and  $2_{Cu_2a}$  over their mononuclear counterparts  $1_{Cu_1a}$  and  $2_{Cu_1a}$ , and show that the (CAAC)CuOTf complex adopts the dinuclear pathway after an initiation period. (B) Evolution of the amount of dimetallated triazole  $2_{Cu_2a}$  and free triazole **3** during the early period of the catalytic reaction using  $1_{Cu_2a}$  as catalyst, showing that  $2_{Cu_2a}$  is the resting state of the catalytic cycle. a.u., arbitrary units. (C) Same as (B), but of  $2_{Cu_1a}$  and **3** using  $1_{Cu_1a}$  as catalyst. (D and E)  $Et_3N$ , as a proton scavenger, allows for the rapid formation of  $1_{Cu_2a}$  from (CAAC)CuOTf and phenyl acetylene (D), and consequently shortens the induction period of the CuAAC reaction described in (A) using (CAAC)CuOTf (E).



**Fig. 3. Mechanistic conclusions: Both the mono- and bis-copper pathways are active in the CuAAC reaction, but the latter is kinetically favored.** The protodemetalation is performed by the alkyne, which regenerates the metallated acetylide, thereby excluding the  $\sigma$ -copper acetylide from the preferred catalytic cycle.

although kinetic data demonstrate their involvement. Moreover, when the trifluoromethanesulfonate is used, investigations of the dinuclear pathway permitted the isolation of a never-postulated bis(copper) triazole complex of type  $2_{Cu_2}$ , which is the resting state of the catalytic cycle.

The alkyne serves as the proton source for the demetallation of  $2_{Cu_2}$ , which regenerates the  $\pi,\sigma$ -bis(copper) acetylide of type  $1_{Cu_2}$ , leaving out complexes of type  $1_{Cu_1}$  from the catalytic cycle. These results led to the broader question of whether bis(metallic) complexes of type  $1_{Cu_2}$  are also key species in other copper-catalyzed organic reactions (33).

## MATERIALS AND METHODS

### General

$^1H$  and  $^{13}C$  NMR spectra were recorded on Bruker Avance 300, Jeol ECA 500, and Varian Inova 500 spectrometers. NMR multiplicities are abbreviated as follows: *s* = singlet, *d* = doublet, *t* = triplet, *q* = quartet, *sept* = septet, *m* = multiplet, *br* = broad signal. Chemical shifts are given in parts per million (ppm) and are referenced to  $SiMe_4$  ( $^1H$ ,  $^{13}C$ ) and  $CFCl_3$  ( $^{19}F$ ). All spectra were obtained at 25°C in the solvent indicated. Coupling constants *J* are given in hertz (Hz). Mass spectra were performed at the University of California San Diego Mass Spectrometry Laboratory. Melting points were measured with an electrothermal MEL-TEMP apparatus. Single crystal x-ray diffraction data were collected on a Bruker Apex II CCD (charge-coupled device) detector using Mo- $K\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ). Crystals were selected under

oil, mounted on nylon loops, and then immediately placed in a cold stream of nitrogen. Structures were solved and refined using Olex2 and SHELXTL. The CAAC and 3,3-diphenyl-1-propyne were prepared following literature procedures, whereas all other starting materials were purchased from commercial sources.

### Synthesis of (CAAC)CuCCPh **1<sub>Cu</sub>a**

*n*-BuLi (2.4 mmol, 2.5 M in hexane) was slowly added to a solution of phenyl acetylene (234 mg, 2.3 mmol) in THF (15 ml) at  $-78^{\circ}\text{C}$ . After 30 min, a solution of (CAAC)CuOAc (1.0 g, 2.3 mmol) in THF (10 ml) was added, and the mixture was stirred at ambient temperature for 1 hour. Volatiles were removed under vacuum, and the remaining solid was extracted with benzene ( $3 \times 10$  ml). After removal of benzene under vacuum, the solid was washed with hexane (20 ml). **1<sub>Cu</sub>a** was obtained as a light yellow solid. Yield 88% (970 mg). mp  $152^{\circ}\text{C}$  [decomposition (dec.)].  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ , 500 MHz):  $\delta = 7.48$  (t,  $J = 7.8$  Hz, 1 H, *p*-H), 7.34 (d,  $J = 7.8$  Hz, 2 H, *m*-H), 7.23 (br d,  $J = 7.1$  Hz, 2 H, *o*- $\text{H}_{\text{C}_6\text{H}_5}$ ), 7.14 (br t,  $J = 7.1$  Hz, 2 H, *m*- $\text{H}_{\text{C}_6\text{H}_5}$ ), 7.08 (br t,  $J = 7.1$  Hz, 1 H, *p*- $\text{H}_{\text{C}_6\text{H}_5}$ ), 2.88 (sept,  $J = 6.9$  Hz, 2 H,  $\text{CHMe}_2$ ), 1.98 (s, 2 H,  $\text{CH}_2$ ), 1.96 to 1.87 (m, 2 H,  $\text{CH}_2$ ), 1.86 to 1.74 (m, 2 H,  $\text{CH}_2$ ), 1.39 to 1.30 (m, 18 H,  $\text{CH}_3$ ,  $\text{CHCH}_3$ ), 1.11 (t,  $J = 7.5$  Hz, 6 H,  $\text{CH}_3$ );  $^{13}\text{C}$  NMR ( $\text{CD}_2\text{Cl}_2$ , 125 MHz):  $\delta = 253.5$  ( $\text{C}_{\text{carbene}}$ ), 145.6 ( $\text{C}_q$ ), 135.1 ( $\text{C}_q$ ), 131.8 ( $\text{CH}_{\text{C}_6\text{H}_5}$ ), 129.9 ( $\text{CH}_{\text{Ar}}$ ), 128.1 ( $\text{CH}_{\text{C}_6\text{H}_5}$ ), 127.7 ( $\text{PhCCCu}$ ), 125.7 ( $\text{CH}_{\text{Ar}}$ ), 125.0 ( $\text{CH}_{\text{C}_6\text{H}_5}$ ), 121.9 ( $\text{C}_{\text{q-C}_6\text{H}_5}$ ), 106.3 ( $\text{PhCCCu}$ ), 81.3 ( $\text{C}_q$ ), 63.3 ( $\text{C}_q$ ), 42.8 ( $\text{CH}_2$ ), 31.4 ( $\text{CH}_2$ ), 29.5, 29.4, 27.3, 22.4, 9.9; high-resolution mass spectrometry (HRMS) [electrospray ionization–time-of-flight mass spectrometry (ESI-TOFMS)]: mass/charge ratio ( $m/z$ ), calculated for  $\text{C}_{30}\text{H}_{40}\text{CuNNa}^+$  500.2349, found 500.2355.

### Synthesis of bis(copper) acetylide complexes **1<sub>Cu2</sub>a**

(CAAC)CuOTf (530 mg, 1 mmol) was added to a solution of **1<sub>Cu</sub>a** (480 mg, 1 mmol) in methylene chloride (20 ml). The reaction was stirred for 5 min at ambient temperature. After removing the solvent under vacuum, **1<sub>Cu2</sub>a** was obtained as a white solid. Yield 95% (952 mg). mp  $174^{\circ}\text{C}$  (dec.).  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ , 500 MHz):  $\delta = 7.44$  (t,  $J = 7.8$  Hz, 2 H, *p*-H), 7.26 (d,  $J = 7.8$  Hz, 4 H, *m*-H), 7.22 (t,  $J = 7.6$  Hz, 1 H, *p*-H), 7.04 (t,  $J = 7.6$  Hz, 2 H, *m*-H), 6.41 (d,  $J = 7.6$  Hz, 2 H, *o*-H), 2.82 (sept,  $J = 6.9$  Hz, 4 H,  $\text{CHMe}_2$ ), 2.05 (s, 4 H,  $\text{CH}_2$ ), 1.96 to 1.83 (m, 4 H,  $\text{CH}_2$ ), 1.80 to 1.68 (m, 4 H,  $\text{CH}_2$ ), 1.38 (s, 12H,  $\text{CH}_3$ ), 1.29 (d,  $J = 6.9$  Hz, 12 H,  $\text{CHCH}_3$ ), 1.18 to 1.02 (m, 24 H);  $^{13}\text{C}$  NMR ( $\text{CD}_2\text{Cl}_2$ , 125 MHz):  $\delta = 249.4$  ( $\text{C}_{\text{carbene}}$ ), 145.3 ( $\text{C}_{\text{q-Aro}}$ ), 134.7 ( $\text{C}_{\text{q-Aro}}$ ), 132.0 ( $\text{CH}_{\text{Aro}}$ ), 130.4 ( $\text{CH}_{\text{Aro}}$ ), 129.4 ( $\text{CH}_{\text{Aro}}$ ), 128.7 ( $\text{CH}_{\text{Aro}}$ ), 125.4 ( $\text{CH}_{\text{Aro}}$ ), 122.2 ( $\text{PhCCCu}_2$ ), 118.9 ( $\text{C}_{\text{q-Aro}}$ ), 110.8 ( $\text{PhCCCu}_2$ ), 82.5 ( $\text{C}_q$ ), 63.1 ( $\text{C}_q$ ), 42.1 ( $\text{CH}_2$ ), 31.5 ( $\text{CH}_2$ ), 29.4, 27.2, 22.4, 9.9; HRMS (ESI-TOFMS):  $m/z$  calculated for  $\text{C}_{52}\text{H}_{75}\text{Cu}_2\text{N}_2^+$  853.4517, found 853.4507.

### Synthesis of bis(copper) triazole complexes **2<sub>Cu2</sub>a**

Benzyl azide (40.0 mg, 0.30 mmol) was added to a solution of **1<sub>Cu2</sub>a** (250 mg, 0.25 mmol) in methylene chloride (0.5 ml). After 1 hour, diethyl ether (15 ml) was added to induce the precipitation of the product. After filtration, **2<sub>Cu2</sub>a** was isolated as a pale yellow solid. Yield 87% (246 mg). mp  $175^{\circ}\text{C}$  (dec.).  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ , 500 MHz):  $\delta = 7.48$  to 7.38 (m, 2 H, *p*-H), 7.34 to 7.14 (m, 12 H), 6.82 (br, 2 H), 5.11 (s, 2 H,  $\text{CH}_2\text{Ph}$ ), 2.77 (sept,  $J = 6.9$  Hz, 4 H,  $\text{CHMe}_2$ ), 2.01 (s, 2 H,  $\text{CH}_2$ ), 1.96 (s, 2 H,  $\text{CH}_2$ ), 1.85 to 1.73 (m, 2 H,  $\text{CH}_2$ ), 1.73 to 1.61 (m, 4 H,  $\text{CH}_2$ ), 1.61 to 1.51 (m, 2 H,  $\text{CH}_2$ ), 1.34 (br, 12 H,  $\text{CH}_3$ ), 1.29 to 1.20 (m, 12 H,  $\text{CHCH}_3$ ), 1.04 to 0.86 (m, 24 H);  $^{13}\text{C}$  NMR ( $\text{CD}_2\text{Cl}_2$ , 125 MHz):  $\delta = 252.7$  ( $\text{C}_{\text{carbene}}$ ), 249.3 ( $\text{C}_{\text{carbene}}$ ), 156.7 ( $\text{C}_{\text{triazolide}}$ ), 155.1 ( $\text{C}_{\text{triazolide}}$ ), 145.4

( $\text{C}_q$ ), 145.3 ( $\text{C}_q$ ), 136.7 ( $\text{C}_{\text{qAr}}$ ), 135.2 ( $\text{C}_q$ ), 134.8 ( $\text{C}_q$ ), 133.3 ( $\text{C}_{\text{qAr}}$ ), 130.3 ( $\text{CH}_{\text{Ar}}$ ), 130.2 ( $\text{CH}_{\text{Ar}}$ ), 129.1 ( $\text{CH}_{\text{Ar}}$ ), 128.8 ( $\text{CH}_{\text{Ar}}$ ), 128.2 ( $\text{CH}_{\text{Ar}}$ ), 128.1 ( $\text{C}_{\text{q-Ar}}$ ), 128.0 ( $\text{CH}_{\text{Ar}}$ ), 127.1 ( $\text{CH}_{\text{Ar}}$ ), 125.2 ( $\text{CH}_{\text{Ar}}$ ), 125.1 ( $\text{CH}_{\text{Ar}}$ ), 82.3 ( $\text{C}_q$ ), 81.9 ( $\text{C}_q$ ), 63.3 ( $\text{C}_q$ ), 63.0 ( $\text{C}_q$ ), 57.1 ( $\text{CH}_2\text{Ph}$ ), 42.4 ( $\text{CH}_2$ ), 31.3 ( $\text{CH}_2$ ), 29.4, 27.0, 22.3, 9.7. HRMS (ESI-TOFMS):  $m/z$  calculated for  $\text{C}_{59}\text{H}_{82}\text{Cu}_2\text{N}_5^+$  986.5157, found 986.5161.

### Protodemetalation of **2<sub>Cu2</sub>a** leading to the regeneration of **1<sub>Cu2</sub>a**

Phenyl acetylene (15 mg, 0.15 mmol) was added to the solution of **2<sub>Cu2</sub>a** (136 mg, 0.12 mmol) in methylene chloride (5 ml). After stirring for 16 hours at room temperature, the solvent was removed under vacuum. The remaining solid was washed with diethyl ether ( $3 \times 10$  ml) to remove triazole **3**, and after drying, **1<sub>Cu2</sub>a** was obtained as a yellow solid. Yield 91% (108 mg). NMR data are identical to those of a sample prepared as described above.

## SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <http://advances.sciencemag.org/cgi/content/full/1/5/e1500304/DC1>

Supplementary text

Fig. S1.  $^1\text{H}$  NMR of (CAAC)CuOAc in  $\text{CD}_2\text{Cl}_2$ .

Fig. S2.  $^{13}\text{C}$  NMR of (CAAC)CuOAc in  $\text{CD}_2\text{Cl}_2$ .

Fig. S3.  $^1\text{H}$  NMR of (CAAC)CuCl in  $\text{CD}_2\text{Cl}_2$ .

Fig. S4.  $^{13}\text{C}$  NMR of (CAAC)CuCl in  $\text{CD}_2\text{Cl}_2$ .

Fig. S5.  $^1\text{H}$  NMR of (CAAC)CuOTf in  $\text{CDCl}_3$ .

Fig. S6.  $^{13}\text{C}$  NMR of (CAAC)CuOTf in  $\text{CDCl}_3$ .

Fig. S7.  $^1\text{H}$  NMR of **1<sub>Cu</sub>a** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S8.  $^{13}\text{C}$  NMR of **1<sub>Cu</sub>a** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S9.  $^1\text{H}$  NMR of **1<sub>Cu2</sub>a** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S10.  $^{13}\text{C}$  NMR of **1<sub>Cu2</sub>a** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S11.  $^1\text{H}$  NMR of **2<sub>Cu2</sub>a** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S12.  $^{13}\text{C}$  NMR of **2<sub>Cu2</sub>a** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S13.  $^1\text{H}$  NMR of **1<sub>Cu2</sub>b** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S14.  $^{13}\text{C}$  NMR of **1<sub>Cu2</sub>b** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S15.  $^1\text{H}$  NMR of **2<sub>Cu</sub>a** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S16.  $^{13}\text{C}$  NMR of **2<sub>Cu</sub>a** in  $\text{CD}_2\text{Cl}_2$ .

Fig. S17. Evolution of the annulation reaction monitored by  $^1\text{H}$  NMR.

Fig. S18. Evolution of the protodemetalation reaction monitored by  $^1\text{H}$  NMR.

Fig. S19. Evolution of the catalytic reaction between benzyl azide and phenyl acetylene monitored by  $^1\text{H}$  NMR.

Table S1. Crystal data and structure refinement for **1<sub>Cu2</sub>a**.

Table S2. Crystal data and structure refinement for **2<sub>Cu2</sub>a** (X = BF<sub>4</sub>).

Table S3. Crystal data and structure refinement for **1<sub>Cu2</sub>b** (X = BF<sub>4</sub>).

## REFERENCES AND NOTES

- V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless, A stepwise Huisgen cycloaddition process: Copper(I)-catalyzed regioselective "ligation" of azides and terminal alkynes. *Angew. Chem. Int. Ed. Engl.* **41**, 2596–2599 (2002).
- C. W. Tornøe, C. Christensen, M. Meldal, Peptidotriazoles on solid phase: [1,2,3]-Triazoles by regioselective copper(I)-catalyzed 1,3-dipolar cycloadditions of terminal alkynes to azides. *J. Org. Chem.* **67**, 3057–3064 (2002).
- H. C. Kolb, M. G. Finn, K. B. Sharpless, Click chemistry: Diverse chemical function from a few good reactions. *Angew. Chem. Int. Ed. Engl.* **40**, 2004–2021 (2001).
- M. G. Finn, V. V. Fokin, Click chemistry: Function follows form. *Chem. Soc. Rev.* **39**, 1231–1232 (2010).
- R. A. Evans, The rise of azide-alkyne 1,3-dipolar 'click' cycloaddition and its application to polymer science and surface modification. *Aust. J. Chem.* **60**, 384–395 (2007).
- P. L. Golas, K. Matyjaszewski, Marrying click chemistry with polymerization: Expanding the scope of polymeric materials. *Chem. Soc. Rev.* **39**, 1338–1354 (2010).
- Y. Hua, A. H. Flood, Click chemistry generates privileged CH hydrogen-bonding triazoles: The latest addition to anion supramolecular chemistry. *Chem. Soc. Rev.* **39**, 1262–1271 (2010).

8. L. Liang, D. Astruc, The copper(I)-catalyzed alkyne-azide cycloaddition (CuAAC) "click" reaction and its applications. An overview. *Coord. Chem. Rev.* **255**, 2933–2945 (2011).
9. X. P. He, J. Xie, Y. Tang, J. Li, G. R. Chen, CuAAC click chemistry accelerates the discovery of novel chemical scaffolds as promising protein tyrosine phosphatases inhibitors. *Curr. Med. Chem.* **19**, 2399–2405 (2012).
10. C. Nájera, J. M. Sansano, 1,3-Dipolar cycloadditions: Applications to the synthesis of antiviral agents. *Org. Biomol. Chem.* **7**, 4567–4581 (2009).
11. A. H. El-Sagheer, T. Brown, Click chemistry with DNA. *Chem. Soc. Rev.* **39**, 1388–1405 (2010).
12. J. M. Meldal, C. W. Tornøe, Cu-catalyzed azide-alkyne cycloaddition. *Chem. Rev.* **108**, 2952–3015 (2008).
13. P. Thirumurugan, D. Matosiuk, K. Jozwiak, Click chemistry for drug development and diverse chemical-biology applications. *Chem. Rev.* **113**, 4905–4979 (2013).
14. M. Meldal, C. W. Tornøe, Cu-catalyzed azide-alkyne cycloaddition. *Chem. Rev.* **108**, 2952–3015 (2008).
15. R. Berg, B. F. Straub, Advancements in the mechanistic understanding of the copper-catalyzed azide-alkyne cycloaddition. *Beilstein J. Org. Chem.* **9**, 2715–2750 (2013).
16. V. O. Rodionov, V. V. Fokin, M. G. Finn, Mechanism of the ligand-free Cu-catalyzed azide-alkyne cycloaddition reaction. *Angew. Chem. Int. Ed. Engl.* **44**, 2210–2215 (2005).
17. F. Himo, T. Lovell, R. Hilgraf, V. V. Fostovtsev, L. Noodleman, K. B. Sharpless, V. V. Fokin, Copper(I)-catalyzed synthesis of azoles. DFT study predicts unprecedented reactivity and intermediates. *J. Am. Chem. Soc.* **127**, 210–216 (2005).
18. M. Ahlquist, V. V. Fokin, Enhanced reactivity of dinuclear copper(I) acetylides in dipolar cycloadditions. *Organometallics* **26**, 4389–4391 (2007).
19. B. F. Straub,  $\mu$ -Acetylide and  $\mu$ -alkenylidene ligands in "click" triazole syntheses. *Chem. Commun.* **2007**, 3868–3870 (2007).
20. B. T. Worrell, J. A. Malik, V. V. Fokin, Direct evidence of a dinuclear copper intermediate in Cu(I)-catalyzed azide-alkyne cycloadditions. *Science* **340**, 457–460 (2013).
21. V. Lavallo, Y. Canac, C. Präsang, B. Donnadiou, G. Bertrand, Stable cyclic (alkyl)(amino)carbenes as rigid or flexible, bulky, electron-rich ligands for transition-metal catalysts: A quaternary carbon atom makes the difference. *Angew. Chem. Int. Ed. Engl.* **44**, 5705–5709 (2005).
22. M. Melaimi, M. Soleilhavoup, G. Bertrand, Stable cyclic carbenes and related species beyond diaminocarbenes. *Angew. Chem. Int. Ed. Engl.* **49**, 8810–8849 (2010).
23. M. Soleilhavoup, G. Bertrand, Cyclic (alkyl)(amino)carbenes (CAACs): Stable carbenes on the rise. *Acc. Chem. Res.* **48**, 256–266 (2015).
24. G. Ung, J. Rittle, M. Soleilhavoup, G. Bertrand, J. C. Peters, Two-coordinate  $\text{Fe}^0$  and  $\text{Co}^0$  complexes supported by cyclic (alkyl)(amino)carbenes. *Angew. Chem. Int. Ed. Engl.* **53**, 8427–8431 (2014).
25. D. S. Weinberger, M. Melaimi, C. E. Moore, A. L. Rheingold, G. Frenking, P. Jerabek, G. Bertrand, Isolation of neutral mono- and dinuclear gold complexes of cyclic (alkyl)(amino)carbenes. *Angew. Chem. Int. Ed. Engl.* **52**, 8964–8967 (2013).
26. C. D. Martin, M. Soleilhavoup, G. Bertrand, Carbene-stabilized main group radicals and radical ions. *Chem. Sci.* **4**, 3020–3030 (2013).
27. Most copper acetylides are polymeric in nature (29) with the exception of a  $(\text{NHC})\text{CuC}\equiv\text{CPh}$  complex (16).
28. I. A. Garbuzova, I. R. Gol'ding, A. N. Schegolikhin, Peculiar intensity distribution in the Raman spectra of copper organoacetylides. *J. Raman Spectrosc.* **26**, 391–395 (1995).
29. C. Nolte, P. Mayer, B. F. Straub, Isolation of a copper(I) triazolide: A "click" intermediate. *Angew. Chem. Int. Ed. Engl.* **46**, 2101–2103 (2007).
30. One of the reviewers suggested that traces of copper(I) dirt could be responsible for the monocopper cycloaddition, a hypothesis that cannot be totally ruled out. Consequently, the rate acceleration can only be greater than 94.
31. R. Bai, G. Zhang, H. Yi, Z. Huang, X. Qi, C. Liu, J. T. Miller, A. J. Kropf, E. E. Bunel, Y. Lan, A. Lei, Cu(II)-Cu(I) synergistic cooperation to lead the alkyne C-H activation. *J. Am. Chem. Soc.* **136**, 16760–16763 (2014).
32. O. Back, M. Henry-Ellinger, C. D. Martin, D. Martin, G. Bertrand,  $^{31}\text{P}$  NMR chemical shifts of carbene-phosphinidene adducts as an indicator of the  $\pi$ -accepting properties of carbenes. *Angew. Chem. Int. Ed. Engl.* **52**, 2939–2943 (2013).
33. S. E. Allen, R. R. Walvoord, R. Padilla-Salinas, M. C. Kozlowski, Aerobic copper-catalyzed organic reactions. *Chem. Rev.* **113**, 6234–6458 (2013).

**Acknowledgments:** We are grateful to the U.S. Department of Energy (DE-FG02-13ER16370) for financial support of this work. Metrical data for the solid-state structures of **1Cu2b**, **1Cu2a**, and **2Cu2a** are available free of charge from the Cambridge Crystallographic Data Centre under reference numbers CCDC-1042482, CCDC-1042483, and CCDC-1042484, respectively. **Competing interests:** The authors declare that they have no competing interests.

Submitted 12 March 2015

Accepted 5 April 2015

Published 12 June 2015

10.1126/sciadv.1500304

**Citation:** L. Jin, D. R. Tolentino, M. Melaimi, G. Bertrand, Isolation of bis(copper) key intermediates in Cu-catalyzed azide-alkyne "click reaction". *Sci. Adv.* **1**, e1500304 (2015).

## Isolation of bis(copper) key intermediates in Cu-catalyzed azide-alkyne "click reaction"

Liqun Jin, Daniel R. Tolentino, Mohand Melaimi and Guy Bertrand

*Sci Adv* 1 (5), e1500304.

DOI: 10.1126/sciadv.1500304

### ARTICLE TOOLS

<http://advances.sciencemag.org/content/1/5/e1500304>

### SUPPLEMENTARY MATERIALS

<http://advances.sciencemag.org/content/suppl/2015/06/09/1.5.e1500304.DC1>

### REFERENCES

This article cites 31 articles, 1 of which you can access for free  
<http://advances.sciencemag.org/content/1/5/e1500304#BIBL>

### PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

---

*Science Advances* (ISSN 2375-2548) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science Advances* is a registered trademark of AAAS.