

## PLANT SCIENCE

# Monolignol ferulate conjugates are naturally incorporated into plant lignins

Steven D. Karlen,<sup>1,2</sup> Chengcheng Zhang,<sup>3</sup> Matthew L. Peck,<sup>3</sup> Rebecca A. Smith,<sup>1,2</sup> Dharshana Padmakshan,<sup>1</sup> Kate E. Helmich,<sup>1,2</sup> Heather C. A. Free,<sup>4,5</sup> Seonghee Lee,<sup>6</sup> Bronwen G. Smith,<sup>5</sup> Fachuang Lu,<sup>1,2</sup> John C. Sedbrook,<sup>7</sup> Richard Sibout,<sup>8</sup> John H. Grabber,<sup>9</sup> Troy M. Runge,<sup>1,10</sup> Kirankumar S. Mysore,<sup>11</sup> Philip J. Harris,<sup>4</sup> Laura E. Bartley,<sup>3\*</sup> John Ralph<sup>1,2\*</sup>

Angiosperms represent most of the terrestrial plants and are the primary research focus for the conversion of biomass to liquid fuels and coproducts. Lignin limits our access to fibers and represents a large fraction of the chemical energy stored in plant cell walls. Recently, the incorporation of monolignol ferulates into lignin polymers was accomplished via the engineering of an exotic transferase into commercially relevant poplar. We report that various angiosperm species might have convergently evolved to natively produce lignins that incorporate monolignol ferulate conjugates. We show that this activity may be accomplished by a BAHD feruloyl-coenzyme A monolignol transferase, *OsFMT1* (AT5), in rice and its orthologs in other monocots.

## INTRODUCTION

One of the major adaptations of terrestrial plants to life on land is their ability to produce lignin for structural strength and defense (1). Stochastically synthesized through stepwise radical coupling of 4-hydroxycinnamyl alcohols (called monolignols, primarily coniferyl and sinapyl alcohols), lignin is a polymer with aryl ether and various other C–O–C and C–C interunit connections (2, 3). Fragmentation of lignin allows polysaccharidase enzymes to access and convert cell wall polysaccharides to monomeric sugars or facilitates cell wall deconstruction to cellulose, hemicelluloses, and lignin fragments. These components are used in economically important processes, including production of paper and other fiber products, second-generation biofuels, and other bioproducts (4, 5). Lignin fragmentation often requires high temperatures and/or harsh chemical treatments to cleave even its weakest interunit bonds (6, 7). However, if chemically labile ester bonds are introduced into the lignin polymer, as can be accomplished by augmenting the prototypical monomers with monolignol ferulate (ML-FA) conjugates (Fig. 1A), then lignin fragmentation can occur under mild pretreatment conditions (Fig. 1B) (8–10). The findings here provide evidence that “zip-lignins,” lignins derived, in part, from ML-FAs, have developed naturally via convergent evolution in diverse angiosperm lineages (Fig. 2).

Initial work on the incorporation of ML-FAs into plant lignin was carried out by engineering poplar trees (*Populus alba* × *Populus grandidentata*) to express a gene from Chinese angelica [*Angelica sinensis* (*As*), a dicotyledonous Chinese medicinal plant] encoding a feruloyl-coenzyme A (CoA) monolignol transferase (*AsFMT*) (8). The *AsFMT*

enzyme couples monolignols with feruloyl-CoA, an intermediate in the monolignol biosynthetic pathway (fig. S1), to produce ML-FAs. *AsFMT* is a member of a family of proteins found in plants and fungi termed BAHD acyltransferases (AT; Fig. 3) (11). *AsFMT*-expressing poplar trees produce ML-FAs and use them in lignification, resulting in improved cell wall saccharification following mild base pretreatment (8).

The lignin assay, derivatization followed by reductive cleavage (DFRC) (12), was used to confirm that ML-FAs were integrally incorporated into the lignin of *AsFMT*-expressing poplar trees (8). DFRC cleaves β-ether bonds to release lignin fragments (12), while leaving the ester linkages of incorporated ML-FAs intact. Hence, DFRC-releasable coniferyl and sinapyl dihydroferulate diacetates (ML-DHFAs; Fig. 1C) are diagnostic for the ML-FAs incorporated into lignin and their release level is proportional to the amount of ML-FAs in the lignin (8). However, because DFRC releases just a fraction of the incorporated ML-FAs as ML-DHFAs, only the relative level of conjugates in the lignin can be determined (8). The threshold to detect DFRC-released ML-DHFAs by gas chromatography–multiple reaction monitoring–mass spectrometry (GC-MRM-MS; table S1) is ~0.01 mg/g of acetyl bromide soluble lignin (ABSL) (see Materials and Methods). These experimental conditions revealed that wild-type (WT) poplar trees already release low levels (0.3 mg/g of ABSL) of the ML-DHFAs from their lignins (8), indicating that poplar plants naturally synthesize ML-FA conjugates and use them in lignification.

## RESULTS

### Plants accumulating ML-FAs in extractives also use them for lignification


Armed with the new knowledge that plants naturally incorporate ML-FAs and an analytical method to diagnostically detect their incorporation into the polymer with suitable sensitivity, we reexamined the lignin of three plants known to produce ML-FAs in their extractives: Chinese angelica (*A. sinensis*, family Apiaceae), kenaf (*Hibiscus cannabinus*, family Malvaceae), and balsa (*Ochroma pyramidale*, family Malvaceae). Performed on isolated lignins, the DFRC assay showed that these plants also used ML-FAs in their lignification (table S2).

### Phylogeny of plants using ML-FAs in lignification

The abovementioned discovery prompted a survey to determine whether the utilization of ML-FAs in lignification was a trait of eudicots or all

<sup>1</sup>Department of Energy Great Lakes Bioenergy Research Center, Wisconsin Energy Institute, University of Wisconsin–Madison, Madison, WI 53726, USA. <sup>2</sup>Department of Biochemistry, University of Wisconsin–Madison, Madison, WI 53706, USA. <sup>3</sup>Department of Microbiology and Plant Biology, The University of Oklahoma, Norman, OK 73019, USA. <sup>4</sup>School of Biological Sciences, The University of Auckland, Auckland, New Zealand. <sup>5</sup>School of Chemical Sciences, The University of Auckland, Auckland, New Zealand. <sup>6</sup>Department of Horticultural Science, IFAS (Institute of Food and Agricultural Sciences) Gulf Coast Research and Education Center, University of Florida, 14625 County Road 672, Wimauma, FL 33598, USA. <sup>7</sup>Department of Energy Great Lakes Bioenergy Research Center, School of Biological Sciences, Illinois State University, Normal, IL 61790, USA. <sup>8</sup>Institut National de la Recherche Agronomique and AgroParisTech, Institut Jean-Pierre Bourgin UMR 1318, Saclay Plant Science, 78000 Versailles, France. <sup>9</sup>U.S. Dairy Forage Research Center, Agricultural Research Service, U.S. Department of Agriculture, 1925 Linden Drive West, Madison, WI 53706, USA. <sup>10</sup>Department of Biological Systems Engineering, University of Wisconsin–Madison, Madison, WI 53706, USA. <sup>11</sup>Plant Biology Division, The Samuel Roberts Noble Foundation, Ardmore, OK 73401, USA.

\*Corresponding author. Email: lbartley@ou.edu (L.E.B.); jralph@wisc.edu (J.R.)



**Fig. 1. Incorporation of ML-FAs into lignin introduces chemically labile esters into the polymer backbone.** (A) FMT enzyme couples feruloyl-CoA and monolignols together to form ML-FA conjugates. The compounds are then transported to the cell wall and undergo radical coupling-based polymerization to form lignin; all the bonds that can be formed when ML-FAs are incorporated into  $\beta$ -ether structures in zip-lignin are shown with dashed lines. (B) Mild base (for example, 0.05 M NaOH at 30°C) cleaves the ML-FA-derived (green) ester bonds dividing the polymer into  $\leq(n + 1)$  fragments, where  $n$  is the number of ML-FA units. (C) DFRC breaks down the lignin by cleaving  $\beta$ -aryl ethers but leaving the esters intact. (D) Electron impact MS fragmentation pattern for coniferyl and sinapyl DHFA (G-DHFA and S-DHFA). FW, formula weight;  $m/z$ , mass/charge ratio. (E) GC-MRM-MS chromatograms of the DFRC product mix reveal the presence of the diagnostic products for ML-FA incorporation into lignin from a number of WT plants. The symbol  $\blacklozenge$  indicates the signals corresponding to S-DHpCA, which shares an MRM transition with G-DHFA.

angiosperms or whether it was a trait ubiquitous to all lignifying plants. We performed the DFRC assay on a set of plants representing the spermatophytes or “seed plants,” including 13 gymnosperms and 54 angiosperms (Fig. 2A). The assayed gymnosperms showed no evidence of ML-FAs, whereas low levels were present in the lignin of many, but not all, of the angiosperms (Fig. 2B and tables S3 to S5). Some of the assayed eudicots showed detectable levels of the diagnostic ML-DHFAs, predominantly derived from coniferyl ferulate (G-FA). Many of the monocots, especially the recently evolved commelinids, which include the major cereal crop plants, tested positive for ML-FAs, with sinapyl ferulate (S-FA) as the main conjugate.


### Origin of ML-FAs and transferase specificity

The discovery of ML-FA production in this diversity of plant species, and notably in plants that use other types of monolignol conjugates for lignification, prompted us to investigate whether ML-FAs arise from a lack of specificity in other transferases or from dedicated FMT enzymes. The “non-specific transferases” hypothesis is most readily addressed in grasses, namely, in rice [*Oryza sativa* (*Os*)] and *Brachypodium* [*Brachypodium distachyon*

(*Bd*)], in which *p*-coumaroyl-CoA monolignol transferases (PMTs), as well as BAHD ATs, have already been identified [*AT4* = *OsPMT1* (13); *Bradi2g36910* = *BdPMT1* (14); and *Bradi2g36980* = *BdPMT2* (15)].

To test the specificity of PMT, we compared the amount of DFRC-releasable ML-DHFAs between a *Brachypodium* line with native *BdPMT1* expression and a sodium azide-generated *BdPmt* missense mutant line that produced lignins devoid of *p*-coumarates (14). Levels of DFRC-released ML-DHFA were similar between the *Brachypodium* WT (0.14 mg/g of ABSL) and mutant lines (0.15 mg/g of ABSL) (Fig. 4 and table S6), indicating that *BdPMT1* activity does not influence ML-FA production and, thus, ML-FAs cannot be attributed to the low specificity of *BdPMT1*.

Furthermore, introduction of the *OsPMT1* gene from rice into the eudicot *Arabidopsis* (*Arabidopsis thaliana*), which does not natively incorporate detectable levels of monolignol conjugates of any kind into its lignin (<0.01 mg/g of ABSL), resulted in transformants with *p*-coumarates esterified to lignin at a level that was easily quantified by DFRC through the release of monolignol dihydro-*p*-coumarates (ML-DHpCAs) (16). However, both the *OsPMT1*-expressing and WT *Arabidopsis* lines did not produce any



**Fig. 2. Comparison of the DFRC-releasable ML-DHFA conjugates among plant species.** (A) A phylogenetic tree of the spermatophytes ("seed plants"), with the orders and families in which plant species were studied. (B) DFRC-released ML-DHFA conjugates; red bars indicate no evidence of ML-DHFAs. Bars indicate SEM for the summation of detected conjugates on duplicate analyses run on a single sample prepared from each plant species.


detectable ML-DHFAs (Fig. 4 and table S6). These two experiments strongly suggest that ML-FA conjugates incorporated into commelinid monocol lignins are not the result of low PMT specificity.

### Identification of a putative new FMT in commelinids

To test the hypothesis that ML-FAs are the product of dedicated FMT enzymes, we examined the annotated genomes of members of the grass family Poaceae, in which ML-FAs appear to be ubiquitous, for proteins with close homology to AsFMT and OsPMT1. We find that these pro-

teins are in distinct BAHD subclades (Fig. 3, clades III and V, respectively). Although rice and other commelinids lack homologs of AsFMT (fig. S2A), they have multiple genes encoding proteins with similarity to PMT (fig. S2B) (17). From screening of rice BAHD mutant lines, we previously reported the preliminary observation that an activation-tagged rice mutant genotype, *OsAT5-D1*, has increased ferulic acid in cell walls compared to WT individuals (17). *OsAT5* is among the most similar proteins to *OsPMT1* (56% identity) and thus seemed to be a reasonable candidate for an enzyme having FMT activity.

Downloaded from <http://advances.sciencemag.org/> on November 18, 2019




Species codes for locus identifiers with order classifications in parentheses are as follows: AT, *Arabidopsis thaliana* (Brassicales); Bradi, *Brachypodium distachyon* (Poales); Eucgr, *Eucalyptus grandis* (Myrtales); Glyma, *Glycine max* (Fabales); GSMUA, *Musa acuminata* (Zingiberales); GRMZM, *Zea mays* (Poales); LOC\_Os, *O. sativa* (Poales); Medtr, *Medicago truncatula* (Fabales); Pavir, *Panicum virgatum* (Poales); PDK, *Phoenix dactylifera* (Areciales); POTR, *Populus trichocarpa* (Malpighiales); and Sb, *Sorghum bicolor* (Poales).

**Fig. 3. Phylogenetic reconstruction of BAHD acyl-CoA ATs is consistent with the convergent evolution of the two feruloyl-CoA monoglignol transferases, OsAT5/FMT and AsFMT.** Maximum likelihood phylogeny of AsFMT, OsAT5, and biochemically characterized BAHD proteins (17). Branch values are based on 1000 bootstraps. Protein IDs are National Center for Biotechnology Information GenBank identifiers or genome locus identifiers.

Here, we provide molecular genetic evidence consistent with the hypothesis that OsAT5 is an FMT. The OsAT5-D1 line has a transfer DNA (T-DNA) insertion with multiple copies of a 35S transcription enhancer upstream of the OsAT5 transcriptional start site (fig. S3A). We confirmed the increased expression of OsAT5 in rice plants with the T-DNA insertion compared with negative segregants lacking the insertion (fig. S3B). In contrast, the expression of more distant genes flanking the insertion site remained unchanged (fig. S3). As hypothesized, OsAT5-D1 plants exhibited an approximately fivefold increase

in the DFRC-releasable ML-DHFA but no increase in ML-DH<sub>p</sub>CA; by comparison, the best AsFMT poplar lines showed an approximately sevenfold increase in released ML-DHFAs (Fig. 4B and table S6) (8). The increase in ferulate is not associated with arabinoxylans, a well-known location of ferulate incorporation in grass cell walls (18), because ferulate released by mild acid hydrolysis remained unchanged (fig. S5). To confirm the association between increased expression of OsAT5 and increased lignin-associated ferulate esters, we generated and characterized three additional overexpression lines, in which OsAT5 expression





**Fig. 4. The amounts of DFRC-releasable ML-DHFA conjugates correlate with the expression of FMT genes but not with the expression of PMT.** (A) No significant change was observed between WT *Brachypodium* and a *Bdpm1* mutant with no PMT activity. Introduction of *BdPMT* into *Arabidopsis* results in detectable ML-DHpCA but no detectable ML-DHFA. (B) Rice overexpressing *OsAT5* (*OsFMT1*), either via activation-tagging in *OsAT5-D1* or via a *Ubi* promoter, and transgenic *AsFMT* poplar show an increase (five- to sevenfold) in ML-DHFAs. Bars indicate SEM of three to seven biological replicates that were measured with technical replicates for each. \* $P < 0.05$ , \*\* $0.01 < P < 0.01$ , and \*\*\* $P < 0.001$ , Student's *t* test.

was driven by the *ZmUbi* promoter (Fig. 4B and fig. S4). All showed increased DFRC-releasable ML-DHFA (Fig. 4B and table S6), compared to the negative segregant controls, establishing that *OsAT5* acts on the monolignols. By all indications, then, *OsAT5* is a native FMT enzyme in rice, that is, *OsAT5* = *OsFMT1*.

Having identified *OsAT5* as a potential grass *FMT* gene, we compared the *OsAT5* sequence to known BAHD AT gene sequences in other commelinid monocots. BAHD proteins were previously identified in the annotations of several plant species (17), and here, we add the data from the draft genomes of banana and date palm (both nongrass commelinids), maize, switchgrass, and eucalyptus. All commelinids examined have several *OsAT5*/*FMT* homologs, whereas this subgroup of ATs has only one or two homologs in the eudicots (fig. S2B). The presence of *OsAT5*-encoding genes corresponds with detectable levels of DFRC-releasable ML-DHFAs, suggesting that the commelinid monocot *FMT* genes may have evolved as a trait of this group.

## DISCUSSION

Upon examination of the phylogeny of the assessed plant species (Fig. 2), some patterns emerged. For example, plants known to produce significant levels of other monolignol conjugates also incorporate ML-FAs into their lignins. These include the commelinids, which incorporate monolignol *p*-coumarates into their lignins and also have ferulates acylating the arabinoxylan hemicelluloses (18), and plants that use monolignol *p*-hydroxybenzoates in lignification, such as palms (*Arecaceae*), willows (*Salix*), and poplars/aspens (*Populus*). A weaker link exists between plants incorporating ML-FAs and those known to use monolignol acetates, which include kenaf and sisal (19), several other monocots (20–22), and various hardwoods (23, 24). However, the introduction of ML-FAs into *Arabidopsis* via transgenesis, which appears to natively lack them, indicates that the ability to use monolignol conjugates does not appear to be isolated to those species that have evolved to do so, highlighting the plasticity of cell wall lignification.

The evidence for BAHD proteins with FMT activity (fig. S2) suggests that the activity has arisen at least twice; that is, it has convergently evolved. For example, the two nonhomologous *FMT* protein sequences, *OsFMT* from a commelinid monocot and *AsFMT* from a eudicot, have little sequence similarity (20%). Furthermore, the model that shows that FMT activity is an ancestral trait of angiosperms is inconsistent with the apparent absence of DFRC-releasable ML-DHFAs in most of the non-commelinid monocots and many of the eudicots, although it cannot be ruled out. Whether an *AsFMT* ortholog is responsible for the incorporation of ML-FAs into eudicot lignins is unclear from the phylogenetic reconstructions. Analysis of the genomes of the ML-FA-producing eudicot tree species *Eucalyptus globulus* and *Populus trichocarpa* reveals a large number of clade III BAHD ATs that are absent from the commelinids, none of which have great similarity to *AsFMT*, as the closest poplar homologs are POPTR\_0001s31750 and POPTR\_0004s01720, with 36% identity (fig. S2A).

The convergent evolution and subsequent proliferation of plants that incorporate ML-FA conjugates into their lignins indicate that, potentially, there is a biological advantage for the production of this lignin structure. Regardless of the actual driving forces selecting for them, the diversity and environmental success of plants with native zip-lignins show that they have no apparent general disadvantages in terms of plant defense or structural stability.

Finally, our findings further refute the contention by some researchers that lignins are derived only from three monolignols. It has been increasingly evident over the past 20 years that many other compounds biosynthesized by plants are used as monomers in lignification (2, 3, 25, 26). As demonstrated here, ML-FA conjugates must now be added to the list of authentic lignin precursors. In practical terms, our discovery unveils new approaches to increasing levels of readily cleavable ester bonds in the lignin backbone, either by breeding or by transgenic methods similar to those used to introduce *AsFMT* into poplar (8). Further work is also needed to explore the effects of ML-FA-containing lignins on processes such as carbon sequestration and biomass utilization.

**MATERIALS AND METHODS****DFRC procedure for ML-DHFA conjugates released from cell wall lignins**

Incorporation of ML-FAs into the lignin was determined using the ether-cleaving, ester-retaining DFRC method previously established for ML-*p*CA and ML-FA conjugates (8, 14, 27, 28). Extract-free cell wall samples (50 mg) or enzyme lignin samples (50 mg) were stirred in 2-dram vials fitted with polytetrafluoroethylene pressure release caps with acetyl bromide/acetic acid [1:4 (v/v), 4 ml]. After heating for 3 hours at 50°C, the solvents were removed by SpeedVac (50°C, 35 min, 1.0 torr, 35 torr/min; Thermo Scientific SPD131DDA). Crude films were suspended in absolute ethanol (0.5 ml), dried on SpeedVac (50°C, 15 min, 6.0 torr, 35 torr/min), and then suspended in dioxane/acetic acid/water [5:4:1 (v/v), 5 ml] with nanopowder zinc (250 mg). The vials were then sealed, sonicated to ensure suspension of solids, and stirred in the dark at room temperature for 16 to 20 hours. Additional nanopowder zinc was added as required to maintain a fine suspension of zinc in the reaction mixtures. The reaction mixtures were then quantitatively transferred with dichloromethane (DCM; 6 ml) into separatory funnels charged with a saturated ammonium chloride (10 ml) and an internal standard (diethyl 5,5'-diferulate diacetate, 54.0 μg). Organics were extracted with DCM (4 × 10 ml), combined, dried over anhydrous sodium sulfate, and filtered, and the solvents were removed via rotary evaporation (water bath at <50°C). Free hydroxyl groups on DFRC products were then acetylated for 16 hours in the dark using a solution of pyridine and acetic anhydride [1:1 (v/v), 5 ml], after which the solvents were removed via a rotary evaporator to yield crude oily films.

To remove most of the polysaccharide-derived products, acetylated DFRC products were loaded onto solid phase extraction (SPE) cartridges (Supelco Supelclean LC-Si SPE tube, 3 ml, product no. 505048) with DCM (2 × 1.0 ml). After elution with hexanes/ethyl acetate [1:1 (v/v), 8 ml], solvents were removed by rotary evaporation, and the products were transferred in stages with GC-MS-grade DCM to final sample volumes of 200 μl into GC-MS vials containing a 300-μl insert. Samples were analyzed on a triple-quadrupole GC-MS/MS (Shimadzu GCMS-TQ8030) operating in MRM mode using synthetic standards for authentication and calibration. The GC program and acquisition parameters are listed in table S1, and the result of the DFRC assay is listed in tables S2 to S4.

**Detection threshold for DFRC-released ML-DHFAs**

When DFRC products are analyzed via a triple-quadrupole GC-MS, operating in MRM mode, the detectable amount of releasable ML-DHFA is ~0.01 mg/g of ABSL. This estimated threshold of detection is partially based on instrument limitations but is mainly limited by the presence of a large amount of polysaccharides, lignin fragments, and the DFRC-released monolignols that create a complex matrix from which the desired ML-DHFA products need to be extracted. Scaling up the DFRC reaction increases not only the amount of ML-DHFA products formed but also the amount of the other matrix components. We have found that the crossover point for improved sensitivity with increasing biomass is ~50 to 100 mg of plant cell walls.

**Estimating the ML-FA incorporation levels**

Most ML-FAs incorporated into lignin form structures that are not cleaved by DFRC and are not released as measurable ML-DHFAs products. Therefore, biomimetically lignified cell walls prepared with known amounts of ML-FAs were subjected to DFRC to estimate the efficiency of ML-DHFA production from ML-FAs incorporated into plant lig-

nins. For this purpose, primary maize cell walls containing bound native peroxidases were lignified by the dropwise addition of separate solutions of dilute hydrogen peroxide and a 1:1 mixture of coniferyl and sinapyl alcohols substituted with either G-FA [0, 8.4, 15.4, and 26.7 weight % (wt%)] or S-FA (0, 9.0, 16.5, and 28.4 wt%) (10). The artificially lignified cell walls were then analyzed by DFRC, and the results were plotted as the weight % of ML-FA used to prepare the cell walls versus the weight % of ML-DHFA released per gram of ABSL. Regression constants for the released ML-DHFAs were determined by fitting the data to a linear model ( $y = mx$ , where  $y = \text{wt\% ML-FA in cell wall dehydrogenation polymer}$  and  $x = \text{wt\% ML-DHFA released}$ ; G-FA:  $m = 53$ ,  $R^2 = 0.99$ ; S-FA:  $m = 34$ ,  $R^2 = 0.99$ ). These results indicate a direct relationship between DFRC-releasable ML-DH conjugates and the quantity of ML-FA in lignin. However, we stress that this approach provides only a rough means of estimating the quantity of ML-FAs incorporated into native plant lignins.

Applying these results to the yields of ML-DHFAs from surveyed plant species suggests that ML-FA incorporation into lignin averaged 1.8 wt% and ranged from 0.2 to 6.6 wt% (tables S2 to S4). The average weight % of ML-FA in eudicots (2.0 wt%) was slightly higher than the average for the monocots (1.7 wt%). The monocots on average released more S-DHFA than G-DHFA (77:23) as compared to the eudicots that mostly released G-DHFA (S-DHFA/G-DHFA, 1:99).

**Procedure to determine ABSL content**

The ABSL contents were measured in 1-cm quartz cuvettes on a Shimadzu UV-1800 spectrophotometer at  $\lambda = 280 \text{ nm}$  and  $\epsilon_{280} = 20.0$ , as previously described (29, 30).

**Preparation of cell wall and enzyme lignin samples**

Cell wall samples were prepared by one of four methods (A to D) described below.

**Method A:** Gymnosperm and angiosperm samples were freeze-dried and shaker-milled into fine powder (Retsch MM 400). Samples were then solvent-extracted sequentially with water (3 × 45 ml), 80% ethanol (3 × 45 ml), and acetone (2 × 45 ml) by first suspending the sample in solvent, sonicating for 20 min, pelleting by centrifugation (8800g for 20 min; Sorvall Biofuge Primo Centrifuge), and, finally, decanting the supernatant. The extract-free pellet was then dried under vacuum for analysis.

**Method B:** Tissues with lignified walls were identified histochemically [phloroglucinol-HCl and ultraviolet fluorescence microscopy (31)]. The lignified tissues were isolated and ground (ROCKLABS Ltd.) in 3-(*N*-morpholino)propanesulfonic acid-KOH buffer [20 mM (pH 6.8)]. The homogenates were centrifuged (1000g for 10 min); resuspended in buffer; filtered through a nylon mesh (11-μm pore size); washed successively with buffer, ethanol, methanol, and *n*-pentane; and air-dried. Starch granules were detected and removed from the preparations from *Cyperus papyrus*, *Hedychium gardnerianum*, and *Phoenix canariensis*, using the method of Carnachan and Harris (32).

**Method C:** Hardwood eudicot samples were Wiley-milled to pass through a 40-mesh screen, air-dried, and extracted three times with a 9:1 mixture of acetone and water. Three additional extractions with methanol were used when required to remove intensely colored extractives. After air-drying, the samples were again Wiley-milled to pass through a 40-mesh screen.

**Method D:** Samples of mature stems were processed to destarched alcohol-insoluble residues (AIR), following a reported procedure (17).

Enzyme lignins were prepared from ball-milled materials, as previously described (33, 34). Briefly, ball-milled extract-free materials (1 g;

prepared using method A) in 50-ml centrifuge tubes were incubated at 35°C and mixed in a shaker at 225 rpm for 3 days with 40 ml of sodium acetate buffer (pH 5.0) and 40 mg of crude cellulases (CELLULYSIN, EMD Biosciences). After incubation, solids were pelleted by centrifugation (8800g for 20 min; Sorvall Biofuge Primo Centrifuge) and washed with acetate buffer (2 × 40 ml). Washed solids were then treated again with the crude cellulases (40 mg) for 3 days and then pelleted and washed with reverse osmosis water (3 × 40 ml). The resulting pelleted solids were dried on a freeze dryer to yield ~10% of the dry weight of the original ball-milled cell wall material.

### Phylogenetic analysis of BAHD ATs

Phylogenetic analysis was conducted in two stages. First, to determine the evolutionary relationship between OsAT5 and AsFMT proteins, we conducted a maximum likelihood phylogenetic reconstruction of three characterized rice ATs (OsAT5, OsAT4/PMT, and OsAT10) and AsFMT with a set of 46 biochemically characterized BAHD enzymes that had previously been divided into five clades, I to V (11), and three PF02458-containing proteins from fungi and one from moss (*Physcomitrella patens*) to serve as an outgroup. We used MEGA5.2.2 (35) to infer and visualize maximum likelihood phylogenies with the following parameters: amino acid substitutions according to the Jones-Taylor-Thornton model,  $\gamma$  distribution of mutation rate among sites, a distribution shape parameter of 5, and gaps treated by partial deletion, allowing site coverage as low as 95%. From these analyses (Fig. 3), we concluded that AsFMT and OsAT5 are not evolutionarily related. Rather, OsAT5, OsAT4, and OsAT10 are closely related to the MsBanAAT in BAHD clade V and belong to a subclade of BAHD proteins, previously referred to as the “Mitchell clade.” By contrast, AsFMT is a member of clade III.

Second, we separately analyzed clade III and the Mitchell clade across diverse species and focused on the genera represented in the DFRC screening (fig. S2, A and B). We included two nongrass commelinid monocots, banana and palm, because of the expectation that these plants might have a close homolog of the FMT in grasses. To identify putative BAHD ATs from the diverse species, we used HMMER version 3.1 (36) with the hidden Markov model profile for PF02458 from the Pfam database. We searched the following genome annotation sources and versions, which were current at the time of the analysis: maize (*Zea mays*), MaizeGDB version 2; switchgrass (*Panicum virgatum*), Phytozome version 1.1; palm (*Phoenix dactylifera*), Weill Cornell Medicine-Qatar PDK sequence version 3; eucalyptus (*Eucalyptus grandis*), Phytozome version 1.1; and banana (*Musa acuminata*), Banana Genome Hub DH-Pahang v1. For comparison with previous analyses, we also included sequences from the following sources: *Arabidopsis* (*A. thaliana*), The *Arabidopsis* Information Resource version 10; soybean (*Glycine max*), Phytozome version 7.0; *Medicago truncatula*, Mt3.5; sorghum (*Sorghum bicolor*), Phytozome version 7.0; rice (*O. sativa*), Michigan State University version 6.1; and *B. distachyon*, Phytozome version 7.0 (17). Phylogenetic analyses were then conducted with the Pfam domain protein sequences (fig. S2B). We determined the BAHD clade of each predicted protein via comparison with the D’Auria set using Clustal2 (37) and omitted sequences that lacked the region surrounding the highly conserved active-site motif HXXXD. From the maximum likelihood phylogenetic reconstructions for these initial single species (100 bootstraps), we identified proteins most closely related to those in the Mitchell clade and clade III. For clade III, we identified 12 proteins from eucalyptus, 9 from *Arabidopsis*, 1 from palm, and none from rice, maize, or banana, suggesting that clade III is not as widely represented in commelinids as in dicots (fig. S2A). For the Mitchell clade, we identified 22 predicted proteins from banana, 20 from

rice, 17 from maize, 23 from switchgrass, 13 from sorghum, 16 from *Brachypodium*, 11 from palm, and 1 from eucalyptus. This builds on the previous results by providing evidence that this clade of genes not only is more abundant in grasses than in eudicotyledonous plants (17) but also includes numerous members in other commelinid monocots. The four subgroups (a to d) of Mitchell clade “I” (within clade V), which have both grass and banana and/or palm proteins, are consistent with there being at least four members of this gene family in the last common ancestor between the grasses and other commelinids. We then carried out maximum likelihood phylogenetic reconstructions with the commelinid-expanded Mitchell clade and related protein sequences and the dicot-expanded clade III. We used 1000 bootstraps for the Mitchell clade tree, with the outgroup consisting of the *Arabidopsis* spermidine dicoumaroyl transferase and spermidine disinapoyl transferase proteins and a group of closely related enzymes that function in taxol biosynthesis. For the clade III tree, we ran 500 bootstraps, and the outgroup consisted of two clade II proteins.

### Gene expression of OsPMT in *A. thaliana*

The OsPMT gene from rice was introduced into *Arabidopsis* (*A. thaliana*), which does not natively incorporate monolignol conjugates of any kind into its lignin, and targeted to secondary cell wall-forming cells (*proCELLULOSE SYNTHASE 7::OsPMT*) (16). These plant lines successfully produced monolignol *p*-coumarates and incorporated them into their lignins.

### Rice lines overexpressing OsFMT (OsAT5)

Rice plants were grown in a greenhouse in Turface Athletics medium/vermiculite (1:1) mix supplemented three times per week with fertilizer (Jack’s Professional LX 15-5-15 4Ca 2Mg) at temperatures from 29° to 32°C during the day and from 24° to 25°C during the night. After germination in water, 7-day-old seedlings were transplanted to the greenhouse. Natural day lengths of less than 13 hours were supplemented with artificial lighting.

We characterized two classes of OsFMT/OsAT5 overexpression rice mutants, an activation-tagged line and three independent *Ubiquitin1* promoter lines. Depending on the assay, we characterized the second (T<sub>2</sub>) and/or third (T<sub>3</sub>) generation of selfed progeny of activation-tagged homozygous mutant and WT segregant rice line of PFG\_2A-20021, referred to herein as OsAT5-D1. The T<sub>1</sub> generation of this line, which is in the Dongjin cultivar background, was previously reported to have an increase in wall-associated ferulates in the mature leaf sheaths (17). We characterized mutant progeny of 2A-20021.10 and WT segregant progeny of 2A-20021.11. DFRC data are from the T<sub>3</sub> generation (mutant 2A-20021.10.7.45.# and WT 2A-20021.11.2.40.#). Trifluoroacetic acid (TFA) fractionation data are from the T<sub>2</sub> generation (mutant 2A-20021.10.2.# and WT 2A-20021.11.7.#), as are gene expression data. In this notation, the period separates generation identities, and the numbers indicate the number of the parental plant of the analyzed progeny. Genotyping, as previously described by Bartley *et al.* (17), was used to isolate homozygous lines and for spot-checking to confirm future generations.

To generate independent lines overexpressing OsAT5, a pUC-based plasmid containing the full coding sequence for OsAT5 was synthesized *de novo*. We then recombined the gene into the pCAMBIA1300-Ubi-GW-Nos construct (38) to produce pCAMBIA1300-UbiAT5. This binary vector contains a Gateway cassette, flanked by the maize *Ubi1* promoter, the 3'-terminator of nopaline synthase from *Agrobacterium tumefaciens*, and the *Hpt2* gene that confers resistance to hygromycin. The rice cultivar Nipponbare was used for transformation after introducing the overexpression



vector *pCAMBIA1300-UbiAT5* into the *A. tumefaciens* strain EH105. The transformation procedure was similar to the method described by Nishimura *et al.* (39). Briefly, embryogenic calli were obtained from 3-week-old immature embryos of Nipponbare. EH105 suspended at an optical density of 0.1 at 600 nm (approximately  $3 \times 10^6$  colony-forming units/ml) was cocultivated with calli for 90 s and placed on sterile filter paper to remove excess *Agrobacterium*. The infected calli were incubated for 60 hours in the dark and, later, were washed and reincubated on regeneration media for 4 weeks under continuous light (3.5 klux). Regenerated plants were grown in deep plastic containers for 1 week, transferred to pots for growth in the greenhouse under the conditions described above, and genotyped with primers for *Hpt2* and the *Ubi<sub>pro</sub>::AT5* construct (table S7). We characterized the T<sub>1</sub> progeny of *Ubi<sub>pro</sub>::AT5* lines—9, 10, and 17—along with negative segregant WT plants for each line. Several additional lines were set aside because of low T<sub>0</sub> seed set or apparent lack of expression of the transgene, assayed as described below.

*OsAT5* overexpression was assayed using quantitative real-time reverse transcription polymerase chain reaction (figs. S3 and S4), with SYBR green and primers as previously described (table S7) (17, 40, 41). Two reference genes, *Ubg5* and *Cc55*, were used for every sample measured. Gene expression was measured in leaf tissue from vegetatively developing rice plants at approximately the V5 stage. We selected samples from the same leaf and developmental stage, that is, developmentally matched, between controls and mutants for each line. For the *OsAT5-D1* line, samples were T<sub>2</sub> mutant and nontransgenic negative segregants approximately 5 weeks after sowing. For the *Ubi<sub>pro</sub>::AT5* lines, samples were second-generation transgenic (T<sub>1</sub>) and negative segregants lacking the transgene from mutant lines 9, 10, and 17.

### Analysis of carbohydrate-associated ferulic acid of *OsAT5-D1*

We used weak acid fractionation to determine whether the changes in hydroxycinnamate content were associated with matrix polysaccharides or lignin. Destarched AIR was created as previously described (17). For weak acid treatment, 2.5 mg of destarched AIR was mixed with 500  $\mu$ l of either 0.05 M TFA or water, similar to a previously described method (17, 42). Samples were incubated with shaking at 100°C for varying times. At each time point, the supernatant (containing solubilized cell wall material) was separated from the remaining solid by centrifugation, and materials were frozen to stop the reaction. Thawed samples were treated with 2 M NaOH and neutralized with concentrated HCl, *trans*-cinnamic acid was added as an internal standard, and the samples were then ethyl acetate-extracted and analyzed by high-performance liquid chromatography as previously described (17).

### SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <http://advances.sciencemag.org/cgi/content/full/2/10/e1600393/DC1>

Supplementary Text

table S1. Chromatography program and MRM parameters for GC-MS/MS characterization of DFRC product mix.

table S2. Experimental results from the analysis of extract-free whole-cell-wall and enzyme lignin samples of selected eudicots.

table S3. Experimental results from the analysis of extract-free whole-cell-wall samples of gymnosperms, magnoliids, and noncommelinid (early) monocots.

table S4. Experimental results from the analysis of extract-free whole-cell-wall samples of commelinid monocots.

table S5. Experimental results from the analysis of extract-free whole-cell-wall samples of eudicots.

table S6. Experimental results from the analysis of extract-free whole-cell-wall samples of plants generated in the enzyme expression study.

table S7. Primers used in this study.

fig. S1. The monolignol biosynthetic pathway indicating the formation of ML-FAs.

fig. S2. The phylogenetic reconstruction of BAHD acyl-CoA ATs is consistent with the convergent evolution of the two feruloyl-CoA monolignol transferases, *OsAT5/FMT* and *AsFMT*.

fig. S3. Genomic position and gene expression data for the *AT5-D1* rice activation-tagged line.

fig. S4. *OsAT5* expression data and DFRC-released ML-DH<sub>p</sub>CA conjugates from *OsAT5* rice lines.

fig. S5. The cell wall compositional differences in *OsAT5-D1* straw are predominantly due to the (50 mM TFA, 100°C) insoluble fraction.

### REFERENCES AND NOTES

- N. D. Bonawitz, C. Chapple, The genetics of lignin biosynthesis: Connecting genotype to phenotype. *Annu. Rev. Genet.* **44**, 337–363 (2010).
- J. Ralph, K. Lundquist, G. Brunow, F. Lu, H. Kim, P. F. Schatz, J. M. Marita, R. D. Hatfield, S. A. Ralph, J. H. Christensen, W. Boerjan, Lignins: Natural polymers from oxidative coupling of 4-hydroxyphenylpropanoids. *Phytochem. Rev.* **3**, 29–60 (2004).
- W. Boerjan, J. Ralph, M. Baucher, Lignin biosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **54**, 519–546 (2003).
- B. M. Upton, A. M. Kasko, Strategies for the conversion of lignin to high-value polymeric materials: Review and perspective. *Chem. Rev.* **116**, 2275–2306 (2016).
- M. FitzPatrick, P. Champagne, M. F. Cunningham, R. A. Whitney, A biorefinery processing perspective: Treatment of lignocellulosic materials for the production of value-added products. *Bioresour. Technol.* **101**, 8915–8922 (2010).
- A. J. Ragauskas, C. K. Williams, B. H. Davidson, G. Britovsek, J. Cairney, C. A. Eckert, W. J. Frederick Jr., J. P. Hallet, D. J. Leak, C. L. Liotta, J. R. Mielenz, R. Murphy, R. Templer, T. Tschalinski, The path forward for biofuels and biomaterials. *Science* **311**, 484–489 (2006).
- C. Somerville, H. Youngs, C. Taylor, S. C. Davis, S. P. Long, Feedstocks for lignocellulosic biofuels. *Science* **329**, 790–792 (2010).
- C. G. Wilkerson, S. D. Mansfield, F. Lu, S. Withers, J.-Y. Park, S. D. Karlen, E. Gonzales-Vigil, D. Padmakshan, F. Unda, J. Rencoret, J. Ralph, Monolignol ferulate transferase introduces chemically labile linkages into the lignin backbone. *Science* **344**, 90–93 (2014).
- J. Ralph, Hydroxycinnamates in lignification. *Phytochem. Rev.* **9**, 65–83 (2010).
- J. H. Grabber, R. D. Hatfield, F. Lu, J. Ralph, Coniferyl ferulate incorporation into lignin enhances the alkaline delignification and enzymatic degradation of maize cell walls. *Biomacromolecules* **9**, 2510–2516 (2008).
- J. C. D'Auria, Acyltransferases in plants: A good time to be BAHD. *Curr. Opin. Plant Biol.* **9**, 331–340 (2006).
- F. Lu, J. Ralph, The DFRC method for lignin analysis. Part 1. New method for  $\beta$ -aryl ether cleavage: Lignin model studies. *J. Agric. Food Chem.* **45**, 4655–4660 (1997).
- S. Withers, F. Lu, H. Kim, Y. Zhu, J. Ralph, C. G. Wilkerson, Identification of a grass-specific enzyme that acylates monolignols with *p*-coumarate. *J. Biol. Chem.* **287**, 8347–8355 (2012).
- D. L. Petrik, S. D. Karlen, C. L. Cass, D. Padmakshan, F. Lu, S. Liu, P. Le Bris, S. Antelme, N. Santoro, C. G. Wilkerson, R. Sibout, C. Lapiere, J. Ralph, J. C. Sedbrook, *p*-Coumaroyl-CoA: monolignol transferase (PMT) acts specifically in the lignin biosynthetic pathway in *Brachypodium distachyon*. *Plant J.* **77**, 713–726 (2014).
- R. Sibout, P. Le Bris, F. Legée, L. Cézard, H. Renault, C. Lapiere, Structural redesigning Arabidopsis lignins into alkali-soluble lignins through the expression of *p*-coumaroyl-CoA: monolignol transferase PMT. *Plant Physiol.* **170**, 1358–1366 (2016).
- R. A. Smith, E. Gonzales-Vigil, S. D. Karlen, J.-Y. Park, F. Lu, C. G. Wilkerson, L. Samuels, J. Ralph, S. D. Mansfield, Engineering monolignol *p*-coumarate conjugates into poplar and Arabidopsis lignins. *Plant Physiol.* **169**, 2992–3001 (2015).
- L. E. Bartley, M. L. Peck, S.-R. Kim, B. Ebert, C. Manisseri, D. M. Chiniquy, R. Sykes, L. Gao, C. Rautengarten, M. E. Vega-Sánchez, P. I. Benke, P. E. Canlas, P. Cao, S. Brewer, F. Lin, W. L. Smith, X. Zhang, J. D. Keasling, R. E. Jentoff, S. B. Foster, J. Zhou, A. Ziebell, G. An, H. V. Scheller, P. C. Ronald, Overexpression of a BAHD acyltransferase, *OsAt10*, alters rice cell wall hydroxycinnamic acid content and saccharification. *Plant Physiol.* **161**, 1615–1633 (2013).
- P. J. Harris, J. A. K. Trethewey, The distribution of ester-linked ferulic acid in the cell walls of angiosperms. *Phytochem. Rev.* **9**, 19–33 (2010).
- J. C. del Río, G. Marques, J. Rencoret, Á. T. Martínez, A. Gutiérrez, Occurrence of naturally acetylated lignin units. *J. Agric. Food Chem.* **55**, 5461–5468 (2007).
- J. Rencoret, J. Ralph, G. Marques, A. Gutiérrez, Á. T. Martínez, J. C. del Río, Structural characterization of the lignin isolated from coconut (*Cocos nucifera*) coir fibers. *J. Agric. Food Chem.* **61**, 2434–2445 (2013).
- J. C. del Río, J. Rencoret, P. Prinsen, Á. T. Martínez, J. Ralph, A. Gutiérrez, Structural characterization of wheat straw lignin as revealed by analytical pyrolysis, 2D-NMR, and reductive cleavage methods. *J. Agric. Food Chem.* **60**, 5922–5935 (2012).
- J. C. del Río, P. Prinsen, J. Rencoret, L. Nieto, J. Jimenez-Barbero, J. Ralph, Á. T. Martínez, A. Gutiérrez, Structural characterization of the lignin in the cortex and pith of elephant grass (*Pennisetum purpureum*) stems. *J. Agric. Food Chem.* **60**, 3619–3634 (2012).
- K. V. Sarkanen, H.-M. Chang, G. G. Allan, Species variation in lignins. 2. Conifer lignins. *Tappi* **50**, 587–590 (1967).



24. J. Ralph, F. Lu, The DFRC method for lignin analysis. Part 6. A modified method to determine acetate regiochemistry on native and isolated lignins. *J. Agric. Food Chem.* **46**, 4616–4619 (1998).
25. Y. Mottiar, R. Vanholme, W. Boerjan, J. Ralph, S. D. Mansfield, Designer lignins: Harnessing the plasticity of lignification. *Curr. Opin. Biotechnol.* **37**, 190–200 (2016).
26. B. A. Simmons, D. Loqué, J. Ralph, Advances in modifying lignin for enhanced biofuel production. *Curr. Opin. Plant Biol.* **13**, 313–320 (2010).
27. F. Lu, J. Ralph, Detection and determination of *p*-coumaroylated units in lignins. *J. Agric. Food Chem.* **47**, 1988–1992 (1999).
28. F. Lu, J. Ralph, in *Lignin: Structural Analysis, Applications in Biomaterials, and Ecological Significance*, F. Lu, Ed. (Nova Science Publishers Inc., 2014), pp. 27–65.
29. R. D. Hatfield, J. Grabber, J. Ralph, K. Brei, Using the acetyl bromide assay to determine lignin concentrations in herbaceous plants: Some cautionary notes. *J. Agric. Food Chem.* **47**, 628–632 (1999).
30. R. S. Fukushima, R. D. Hatfield, Extraction and isolation of lignin for utilization as a standard to determine lignin concentration using the acetyl bromide spectrophotometric method. *J. Agric. Food Chem.* **49**, 3133–3139 (2001).
31. B. G. Smith, P. J. Harris, Polysaccharide composition of unligified cell-walls of pineapple [*Ananas comosus* (L.) Merr.] fruit. *Plant Physiol.* **107**, 1399–1409 (1995).
32. S. M. Carnachan, P. J. Harris, Polysaccharide compositions of primary cell walls of the palms *Phoenix canariensis* and *Rhopalostylis sapida*. *Plant Physiol. Biochem.* **38**, 699–708 (2000).
33. H.-m. Chang, E. B. Cowling, W. Brown, E. Adler, G. Miksche, Comparative studies on cellulolytic enzyme lignin and milled wood lignin of sweetgum and spruce. *Holzforschung* **29**, 153–159 (1975).
34. A. Wagner, J. Ralph, T. Akiyama, H. Flint, L. Phillips, K. Torr, B. Nanayakkara, L. T. Kiri, Exploring lignification in conifers by silencing hydroxycinnamoyl-CoA:shikimate hydroxycinnamoyltransferase in *Pinus radiata*. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 11856–11861 (2007).
35. K. Tamura, D. Peterson, N. Peterson, G. Stecher, M. Nei, S. Kumar, MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731–2739 (2011).
36. R. D. Finn, J. Clements, S. R. Eddy, HMMER web server: Interactive sequence similarity searching. *Nucleic Acids Res.* **39**, W29–W37 (2011).
37. M. A. Larkin, G. Blackshields, N. P. Brown, P. A. McGettigan, H. McWilliam, F. Valentin, I. M. Wallace, A. Wilm, R. Lopez, J. D. Thompson, T. J. Gibson, D. G. Higgins, Clustal W and clustal X version 2.0. *Bioinformatics* **23**, 2947–2948 (2007).
38. C.-J. Park, R. Bart, M. Chern, P. E. Canlas, W. Bai, P. C. Ronald, Overexpression of the endoplasmic reticulum chaperone BiP3 regulates XA21-mediated innate immunity in rice. *PLOS One* **5**, e9262 (2010).
39. A. Nishimura, I. Aichi, M. Matsuoka, A protocol for *Agrobacterium*-mediated transformation in rice. *Nat. Protoc.* **1**, 2796–2802 (2007).
40. F. Piston, C. Uauy, L. Fu, J. Langston, J. Labavitch, J. Dubcovsky, Down-regulation of four putative arabinoxylan feruloyl transferase genes from family PF02458 reduces ester-linked ferulate content in rice cell walls. *Planta* **231**, 677–691 (2010).
41. M. Jain, Genome-wide identification of novel internal control genes for normalization of gene expression during various stages of development in rice. *Plant Sci.* **176**, 702–706 (2009).
42. L. Saulnier, J. Vigouroux, J.-F. Thibault, Isolation and partial characterization of feruloylated oligosaccharides from maize bran. *Carbohydr. Res.* **272**, 241–253 (1995).
43. S. Ouyang, S. Ouyang, W. Zhu, J. Hamilton, H. Lin, M. Campbell, K. Childs, F. Thibaud-Nissen, R. L. Malek, Y. Lee, L. Zheng, J. Orvis, B. Haas, J. Wortman, C. R. Buell, The TIGR rice genome annotation resource: Improvements and new features. *Nucleic Acids Res.* **35**, D883–D887 (2007).

**Acknowledgments:** We thank the University of Wisconsin–Madison Plant Growth Facilities, J. Mullet, M. Casler, S. Ralph, R. Arango, S. Liu, M. Phillips, V. Phillips, and Dannance Karlen Holsteins for providing the biological samples; S. Thibivilliers, D. Petrik, C. Cass, and B. Ebert for their technical support; and P. Ronald for her comments on the manuscript. **Funding:** S.D.K., R.A.S., D.P., K.E.H., F.L., J.C.S., T.M.R., and J.R. were funded by the Department of Energy (DOE) Great Lakes Bioenergy Research Center [DOE Office of Science Biological and Environmental Research (BER) DE-FC02-07ER64494]. M.L.P., C.Z., and L.E.B. were funded in part by the DOE Feedstock Genomics (DOE Office of Science BER DE-SC006904) and in part by the NSF Experimental Program to Stimulate Competitive Research, under grant no. EPS-0814361. H.C.A.F., B.G.S., and P.J.H. were funded in part by a Specific Cooperative Agreement #58-3655-8-129f with the U.S. Dairy Forage Research Center, U.S. Department of Agriculture–Agricultural Research Service, via funding from the DOE Energy Biosciences Program [DE-AI02-06ER64299 (2006)]; H.C.A.F. also received a doctoral scholarship from The University of Auckland. S.L. and K.S.M. were funded by the Samuel Roberts Noble Foundation. **Author contributions:** S.D.K., P.J.H., B.G.S., J.C.S., J.H.G., T.M.R., L.E.B., and J.R. designed the project. S.D.K., M.L.P., C.Z., R.A.S., D.P., K.E.H., H.C.A.F., S.L., and F.L. processed the samples and analyzed the data. S.L., R.A.S., R.S., and K.S.M. prepared the transgenic plants. S.D.K. and J.H.G. prepared the maize cell wall dehydrogenation polymers. S.D.K., R.A.S., J.H.G., P.J.H., B.G.S., L.E.B., and J.R. wrote the manuscript. All authors discussed the results and commented on the draft. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** The DNA sequences for the *AsFMT* (accession no. KF977606) and *OsFMT* (*OsATS*; accession no. NP\_001047408.1) genes are available in GenBank. The rice acyltransferase *AT5* gene corresponds to *LOC\_Os05g19910* [Michigan State University Rice Genome Annotation version 7 (43)]. All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Additional data related to this paper may be requested from the authors.

Submitted 23 February 2016

Accepted 1 September 2016

Published 14 October 2016

10.1126/sciadv.1600393

**Citation:** S. D. Karlen, C. Zhang, M. L. Peck, R. A. Smith, D. Padmakshan, K. E. Helmich, H. C. A. Free, S. Lee, B. G. Smith, F. Lu, J. C. Sedbrook, R. Sibout, J. H. Grabber, T. M. Runge, K. S. Mysore, P. J. Harris, L. E. Bartley, J. Ralph, Monolignol ferulate conjugates are naturally incorporated into plant lignins. *Sci. Adv.* **2**, e1600393 (2016).

## Monolignol ferulate conjugates are naturally incorporated into plant lignins

Steven D. Karlen, Chengcheng Zhang, Matthew L. Peck, Rebecca A. Smith, Dharshana Padmakshan, Kate E. Helmich, Heather C. A. Free, Seonghee Lee, Bronwen G. Smith, Fachuang Lu, John C. Sedbrook, Richard Sibout, John H. Grabber, Troy M. Runge, Kirankumar S. Mysore, Philip J. Harris, Laura E. Bartley and John Ralph

*Sci Adv* 2 (10), e1600393.  
DOI: 10.1126/sciadv.1600393

### ARTICLE TOOLS

<http://advances.sciencemag.org/content/2/10/e1600393>

### SUPPLEMENTARY MATERIALS

<http://advances.sciencemag.org/content/suppl/2016/10/11/2.10.e1600393.DC1>

### REFERENCES

This article cites 42 articles, 9 of which you can access for free  
<http://advances.sciencemag.org/content/2/10/e1600393#BIBL>

### PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

---

*Science Advances* (ISSN 2375-2548) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science Advances* is a registered trademark of AAAS.

Copyright © 2016, The Authors