

## ECOLOGY

# Comment (2) on “Formation of the Isthmus of Panama” by O’Dea *et al.*

Peter Molnar\*

O’Dea *et al.* challenged the inference that the Isthmus of Panama has been in place for the last 10 million years or more and from “an exhaustive review and reanalysis of geological, paleontological, and molecular records,” they argued for a “formation of the Isthmus of Panama *sensu stricto* around 2.8 Ma.” I review environmental changes since ~5 Ma throughout Earth, and I argue that environmental changes in the Central American-Caribbean region have been part of a concurrent, worldwide phenomenon that requires a global, not local, explanation. Accordingly, evidence of environmental change from the Central American-Caribbean region does not implicate the emergence of the Isthmus of Panama.

## INTRODUCTION

Building on a pair of previous studies (1, 2), Montes *et al.* (3) showed that zircons in fluvial sediment deposited in northern Colombia since at least 10 million years ago (Ma) bore an isotopic signature of granitic rock from Panama. Recognizing that rocks neither swim nor would likely be carried by ice to northern Colombia, they inferred that rivers not only deposited the sediment but also must have transported it from the granitic belt in Panama to the sites of deposition. Most investigators have placed the now-closed Central American Seaway, sometimes called the Bolivar Trough (4–7), Bolivar seaway (8), Bolivar Portal (9), Bolivar geosyncline (10), or Atrato Basin (11) in the region between the sediment and the granitic belt, although some had inferred previously that no connection between the Pacific and Atlantic lay there (12). Concurrent with the work of Montes *et al.*, Bacon *et al.* (13) exploited molecular and fossil constraints on the timing of biotic interchange between North and South America to conclude that interchange was well underway before ~3 to 4 Ma, the commonly assumed time of closure of the Central American Seaway. Against these observations for a relatively old land bridge connecting North and South America, if not necessarily isolating the Caribbean Sea from the Pacific Ocean, O’Dea *et al.* (14) argued that much evidence does require a relatively young isthmus (~3 Ma). Although perhaps biased (15), I contend that even if the isthmus proves to be so young, the arguments of O’Dea *et al.* (14) ought to convince no one of such youth.

First, O’Dea *et al.* (14) addressed little of the evidence that bears on the possibility that the Isthmus was in place at 10 Ma. They stated, “In the 1970s, high-resolution paleoceanographic data available from deep-sea cores began to show that an isthmus...was in place only relatively recently, around 3 million years ago (Ma),” and they supported that claim by citing two papers (16, 17). Readers of that sentence might not realize that those two papers presented data applicable to the periods since 4 and 6 Ma, respectively. Neither could address the possibility that the Central America Seaway closed at 10 Ma or earlier.

Second and more importantly, much evidence shows different regional climates across the globe before and after ~3 to 4 Ma. The Northern Hemisphere before 3 to 4 Ma was warmer than today by several degrees Celsius, as shown by environmental conditions in mid- and high-latitude regions on land in North America (18–20) and in Europe (21, 22), along Atlantic coastal regions of North America (23–25), and

by North Atlantic sea surface temperatures (SSTs) (26–28). The largest changes in SSTs come from regions of marked upwelling at lower latitudes, ~9°C off the coast of southern California (29) and ~4°C of the west coast of Africa in the North Atlantic (30), as well as in the South Atlantic, ~10°C off the southwest coast of Africa (31, 32). Whether these regions, where cold, deep water upwells today, cooled because winds changed or because the thermocline shoaled (33) is not pertinent here, for with these changes in SSTs, winds must also have changed.

Marked changes in winds are virtually certain in the equatorial Pacific because the eastern Pacific has cooled by as much as 3° to 4°C since 4 to 5 Ma (34–37). Although some controversy continues, the western equatorial Pacific does not seem to have cooled since that time. Moreover, the thermocline in the eastern equatorial Pacific has shoaled since 4 Ma, as one would expect from strengthened easterly winds (38, 39). This cooling of the eastern Pacific has transformed the SST distribution from one resembling that during occasional El Niño events to the present-day more La Niña-like normal state (33, 40–43). Whereas most of the world was warmer in Early to Middle Pliocene time than today, one area that seems to have been cooler—the southeastern United States and the region surrounding the Gulf of Mexico (23–25, 41, 42)—is the only area that cools during El Niño events (44, 45).

The El Niño-like SST distribution bears on an observation that O’Dea *et al.* (14) used to infer a closing of the Central American Seaway since 3 to 4 Ma; since that time, salinity in the Caribbean has increased (46–51), while the easternmost Pacific freshened (35). O’Dea *et al.* (14) attributed this increased difference in salinity to the isolating of the Caribbean from the Pacific by the emergence of the Isthmus of Panama. A change from El Niño-like conditions and its effect on moisture transport by the atmosphere offer another possible explanation (52). Virtually, all moisture exported from the Atlantic Ocean leaves the Caribbean and crosses Central America to the Pacific (53). Today, less moisture is transported across Central America during El Niño events than under normal conditions (52, 54). The scatter in data is too large to demonstrate that the El Niño-like eastern Pacific SST conditions made moisture transport at 3 to 4 Ma sufficiently small to account for the change in salinity contrast (52), but the El Niño-like SST distribution at 3 to 4 Ma offers a possible explanation for the smaller salinity contrast at that time than today, a possibility that O’Dea *et al.* (14) ignored.

O’Dea *et al.* (14) attributed another difference between the Caribbean and the Pacific to the emergence of Panama. Today, the annual range of shallow-ocean temperature is only ~2°C on the Caribbean side but ~6°C on the Pacific side. From variations in zooid sizes in fossil bryozoan colonies in the Caribbean and the Pacific, O’Dea *et al.* (55) inferred

Copyright © 2017  
The Authors, some  
rights reserved;  
exclusive licensee  
American Association  
for the Advancement  
of Science. Distributed  
under a Creative  
Commons Attribution  
NonCommercial  
License 4.0 (CC BY-NC).

Downloaded from <http://advances.sciencemag.org/> on June 20, 2019

Department of Geological Sciences, Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, Boulder, CO 80309-0399, USA.

\*Corresponding author. Email: molnar@colorado.edu

a change of ~5°C in the mean annual ranges of temperature (MART) since 3 to 4 Ma. O'Dea *et al.* (14) assigned uncertainties of ~2.5°C for MART in the Caribbean and ~1°C for the Pacific. McClelland *et al.* (56) had assessed the statistics of these data, however, and had written: "The revised equation...has a large margin of error ( $\pm 4^\circ\text{C}$ ) and is therefore useful only for comparing environments with a difference in MART greater than 4°C." With uncertainties of 4°C, the inferred ~5°C difference in MART can, at best, be only barely resolved, and given the other global changes at the same time, these inferred differences in MART do not provide a strong argument for the emergence of the isthmus since 4 Ma.

Changes in regional climates since 3 to 4 Ma characterize more than temperatures, winds, and salinity. Many regions became more arid, such as East Africa (57, 58), Chad (59, 60), West Africa (61), the Namib Desert in southwest Africa (62), the western United States (63, 64), and Australia (65–67). Loess deposition increased in China (68), as did dust transport to the North Pacific (69, 70) and to the Southern Ocean (71). High dust accumulation rates indicate arid source regions (72). As with temperature, however, at least one change of opposite sense seems to have occurred; Hovan (73) inferred that dust transport from South America to the eastern Pacific decreased and that northern South America has become wetter.

Finally, although also controversial (74), abundant evidence suggests that both erosion rates (75, 76) and sedimentation accumulation rates (77–79) increased since ~3 to 4 Ma. In particular, studies of either sediment mass accumulation rates or cooling of rock as it has been exhumed show accelerated erosion since ~3 to 4 Ma in diverse settings: the Mississippi River drainage basin (80, 81), the Colorado Plateau (82), Alaska (83, 84), the Alps (85), the Tien Shan (86), and, in at least, parts of the Himalayas (87–89). Moreover, accelerated deposition of sediment could have affected nutrients positively or negatively, with corresponding changes in productivity.

Readers should see that I do not focus on the veracity or applicability of evidence suggesting that the Isthmus of Panama was in place by 10 Ma, nor do I pretend to show that an emergence of the Isthmus did not occur at 3 to 4 Ma. Rather, I contend that O'Dea *et al.* (14) have ignored abundant evidence showing that much happened throughout Earth at the suggested time when the Isthmus of Panama emerged. The most parsimonious interpretation of these global observations is that the evidence for environmental change in the Central America region is part of that global process and does not require a local cause. Some might argue that the closing of the Central America Region at ~3 Ma not only explains the data that O'Dea *et al.* (14) discussed but also caused the global changes noted above. I addressed that idea elsewhere and found little support for it (15). In summary, I would rewrite the final sentence of their abstract with the words "an older isthmus" changed to "a 3-Ma," "before" changed to "only since," and "we" to "I" so that it reads: "The evidence used to support" a 3-Ma "isthmus is inconclusive, and" I "caution against the uncritical acceptance of an isthmus" only since "the Pliocene."

## REFERENCES AND NOTES

- C. Montes, A. Cardona, R. McFadden, S. E. Morón, C. A. Silva, S. Restrepo-Moreno, D.A. Ramírez, N. Hoyos, J. Wilson, D. Farris, G. A. Bayona, C. A. Jaramillo, V. Valencia, J. Bryan, J. A. Flores, Evidence for middle Eocene and younger land emergence in central Panama: Implications for Isthmus closure. *Geol. Soc. Am. Bull.* **124**, 780–799 (2012).
- C. Montes, G. Bayona, A. Cardona, D. M. Buchs, C. A. Silva, S. Morón, N. Hoyos, D. A. Ramírez, C. A. Jaramillo, V. Valencia, Arc-continent collision and orocline formation: Closing of the Central American seaway. *J. Geophys. Res.* **117**, B04105 (2012).
- C. Montes, A. Cardona, C. A. Jaramillo, A. Pardo, J. C. Silva, V. Valencia, C. Ayala, L. C. Pérez-Angel, L. A. Rodríguez-Parra, V. Ramírez, H. Niño, Middle Miocene closure of the Central American Seaway. *Science* **348**, 226–229 (2015).
- L. G. Marshall, Land Mammals and the Great American Interchange. *Am. Sci.* **76**, 380–388 (1988).
- L. G. Marshall, S. D. Webb, J. J. Sepkoski Jr., D. M. Raup, Mammalian evolution and the great American interchange. *Science* **215**, 1351–1357 (1982).
- F. G. Stehli, S. D. Webb, "A kaleidoscope of plates, faunal and floral dispersals, and sea level changes," in *The Great American Biotic Interchange*, F. G. Stehli, S. D. Webb, Eds. (Plenum Press, 1985), pp. 3–16.
- F. C. Whitmore Jr., R. H. Stewart, Miocene mammals and Central American seaways. *Science* **148**, 180–185 (1965).
- F. B. Van Houten, R. B. Travis, Cenozoic deposits, Upper Magdalena Valley, Colombia. *Am. Assoc. Pet. Geol. Bull.* **52**, 675–702 (1968).
- J. Hanken, D. B. Wake, Genetic differentiation among plethodontid salamanders (genus *Bolitoglossa*) in Central and South America: Implications for the South American invasion. *Herpetologica* **38**, 272–287 (1982).
- H. J. Harrington, Paleogeographic development of South America. *Am. Assoc. Pet. Geol. Bull.* **46**, 1773–1814 (1962).
- H. Duque-Caro, "Structural style, diapirism, and accretionary episodes of the Sinú-San Jacinto terrane, southwestern Caribbean borderland," in *The Caribbean-South America Plate Boundary and Regional Tectonics*, W. E. Bonini, R. B. Hargraves, R. Shagam, Eds. (Geological Society of America Memoir 162, 1984), pp. 303–316.
- A. G. Coates, L. S. Collins, M.-P. Aubry, W. A. Berggren, The geology of the Darien, Panama, and the late Miocene-Pliocene collision of the Panama arc with northwestern South America. *GSA Bull.* **116**, 1327–1344 (2004).
- C. D. Bacon, D. Silvestro, C. Jaramillo, B. T. Smith, P. Chakrabarty, A. Antonelli, Biological evidence supports an early and complex emergence of the Isthmus of Panama. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 6110–6115 (2015).
- A. O'Dea, H. A. Lessios, A. G. Coates, R. I. Eytan, S. A. Restrepo-Moreno, A. L. Cione, L. S. Collins, A. de Queiroz, D. W. Farris, R. D. Norris, R. F. Stallard, M. O. Woodburne, O. Aguilera, M.-P. Aubry, W. A. Berggren, A. F. Budd, M. A. Cozzuol, S. E. Coppard, H. Duque-Caro, S. Finnegan, G. M. Gasparini, E. L. Grossman, K. G. Johnson, L. D. Keigwin, N. Knowlton, E. G. Leigh, J. S. Leonard-Pingel, P. B. Marko, N. D. Penson, P. G. Ravello-Dolmen, E. Soibelzon, L. Soibelzon, J. A. Todd, G. J. Vermeij, J. B. C. Jackson, Formation of the Isthmus of Panama. *Sci. Adv.* **2**, e1600883 (2016).
- P. Molnar, Closing of the Central American Seaway and the ice age: A critical review. *Paleoceanography* **23**, PA2201 (2008).
- L. D. Keigwin Jr., Pliocene closing of the Isthmus of Panama, based on biostratigraphic evidence from nearby Pacific Ocean and Caribbean Sea cores. *Geology* **6**, 630–634 (1978).
- T. Saito, Geologic significance of coiling direction in the planktonic foraminifera *Pulleniatina*. *Geology* **4**, 305–309 (1976).
- A. P. Ballantyne, D. R. Greenwood, J. S. Sinninghe Damsté, A. Z. Csank, J. J. Eberle, N. Rybczynski, Significantly warmer Arctic surface temperatures during the Pliocene indicated by multiple independent proxies. *Geology* **38**, 603–606 (2010).
- J. Brigham-Grette, M. Melles, P. Minyuk, A. Andreev, P. Tarasov, R. DeConto, S. Koenig, N. Nowaczyk, V. Wennrich, P. Rosén, E. Haltia, T. Cook, C. Gebhardt, C. Meyer-Jacob, J. Snyder, U. Herzschuh, Pliocene warmth, polar amplification, and stepped pleistocene cooling recorded in NE Arctic Russia. *Science* **340**, 1421–1427 (2013).
- R. H. Tedford, C. R. Harington, An Arctic mammal fauna from the Early Pliocene of North America. *Nature* **425**, 388–390 (2003).
- S. Fauquette, J.-P. Suc, J. Guiot, F. Diniz, N. Feddi, Z. Zheng, E. Bessais, A. Drivalieri, Climate and biomes in the West Mediterranean area during the Pliocene. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **152**, 15–36 (1999).
- S. Panitz, U. Salzmann, B. Risebrobakken, S. De Schepper, M. J. Pound, Climate variability and long-term expansion of peatlands in Arctic Norway during the late Pliocene (ODP Site 642, Norwegian Sea). *Clim. Past* **12**, 1043–1060 (2016).
- T. M. Cronin, R. Whitley, A. Wood, A. Tsukagoshi, N. Ikeya, E. M. Brouwers, W. M. Briggs Jr., Microfaunal evidence for elevated mid-Pliocene temperatures in the Arctic Ocean. *Paleoceanography* **8**, 161–173 (1993).
- D. A. Willard, Palynological record from the North Atlantic region at 3 Ma: Vegetational distribution during a period of global warmth. *Rev. Paleobot. Palynol.* **83**, 275–297 (1994).
- D. A. Willard, T. M. Cronin, S. E. Ishman, R. J. Litwin, Terrestrial and marine records of climate and environmental change during the Pliocene in subtropical Florida. *Geology* **21**, 679–682 (1993).
- K. T. Lawrence, T. D. Herbert, C. M. Brown, M. E. Raymo, A. M. Haywood, High-amplitude variations in North Atlantic sea surface temperature during the early Pliocene warm period. *Paleoceanography* **24**, PA2218 (2009).
- K. T. Lawrence, S. Sosdian, H. E. White, Y. Rosenthal, North Atlantic climate evolution through the Plio-Pleistocene climate transitions. *Earth Planet. Sci. Lett.* **300**, 329–342 (2010).

28. M. Mudelsee, M. E. Raymo, Slow dynamics of the Northern Hemisphere glaciation. *Paleoceanography* **20**, PA4022 (2005).
29. P. S. Dekens, A. C. Ravelo, M. D. McCarthy, Warm upwelling regions in the Pliocene warm period. *Paleoceanography* **22**, PA3211 (2007).
30. T. D. Herbert, J. D. Schuffert, "Alkenone unsaturation estimates of late Miocene through late Pliocene sea-surface temperatures at Site 958," in *Proc. Ocean Drill. Program Sci. Results 1597*, J. V. Firth, Ed. (Ocean Drilling Program, College Station, 1998), pp. 17–22.
31. J. R. Marlow, C. B. Lange, G. Wefer, A. Rosell-Melé, Upwelling intensification as part of the Pliocene-Pleistocene climate transition. *Science* **290**, 2288–2291 (2000).
32. A. Rosell-Melé, A. Martínez-García, E. L. McClymont, Persistent warmth across the Benguela upwelling system during the Pliocene epoch. *Earth Planet. Sci. Lett.* **386**, 10–20 (2014).
33. S. G. Philander, A. V. Fedorov, Role of tropics in changing the response to Milankovich forcing some three million years ago. *Paleoceanography* **18**, 1045 (2003).
34. W. P. Chaisson, A. C. Ravelo, Pliocene development of the east-west hydrographic gradient in the tropical Pacific. *Paleoceanography* **15**, 497–505 (2000).
35. J. Groeneveld, S. Steph, R. Tiedemann, D. Garbe-Schönberg, D. Nürnberg, A. Sturm, "Pliocene mixed-layer oceanography for Site 1241, using combined Mg/Ca and  $\delta^{18}\text{O}$  analyses of *Globigerinoides sacculifer*," in *Proc. Ocean Drill. Program Sci. Results 202*, R. Tiedemann, A. C. Mix, C. Richter, W. F. Ruddiman, Eds. (Ocean Drilling Program, College Station, 2006), pp. 1–27.
36. K. T. Lawrence, Z. Liu, T. D. Herbert, Evolution of the eastern tropical Pacific through Plio-Pleistocene glaciation. *Science* **312**, 79–83 (2006).
37. M. W. Wara, A. C. Ravelo, M. L. Delaney, Permanent El Niño-like conditions during the Pliocene warm period. *Science* **309**, 758–761 (2005).
38. K. G. Cannariato, A. C. Ravelo, Plio-Pleistocene evolution of eastern tropical Pacific rate of circulation and thermocline depth. *Paleoceanography* **12**, 805–820 (1997).
39. H. L. Ford, A. C. Ravelo, P. S. Dekens, J. P. LaViviere, M. W. Wara, The evolution of the equatorial thermocline and the early Pliocene *El Padre* mean state. *Geophys. Res. Lett.* **42**, 4878–4887 (2015).
40. A. V. Fedorov, P. S. Dekens, M. McCarthy, A. C. Ravelo, P. B. deMenocal, M. Barreiro, R. C. Pacanowski, S. G. Philander, The Pliocene Paradox (Mechanisms for a Permanent El Niño). *Science* **312**, 1485–1489 (2006).
41. P. Molnar, M. A. Cane, El Niño's tropical climate and teleconnections as a blueprint for pre-Ice Age climates. *Paleoceanography* **17**, 11–1–11–11 (2002).
42. P. Molnar, M. A. Cane, Early Pliocene (pre-Ice Age) El Niño-like Global Climate: Which El Niño? *Geosphere* **3**, 337–365 (2007).
43. A. C. Ravelo, P. S. Dekens, M. McCarthy, Evidence for El Niño-like conditions during the Pliocene. *GSA Today* **16**, 4–11 (2006).
44. G. N. Kiladis, H. F. Diaz, An analysis of the 1877–1878 ENSO episode and comparison with 1982–83. *Mon. Weath. Rev.* **114**, 1035–1047 (1986).
45. C. F. Ropelewski, M. S. Halpert, North American precipitation and temperature patterns associated with the El Niño/Southern Oscillation (ENSO). *Mon. Weath. Rev.* **114**, 2352–2362 (1986).
46. J. Groeneveld, D. Nürnberg, R. Tiedemann, G.-J. Reichert, S. Steph, L. Reuning, D. Crudeli, P. Mason, Foraminiferal Mg/Ca increase in the Caribbean during the Pliocene: Western Atlantic Warm Pool formation, salinity influence, or diagenetic overprint? *Geochem. Geophys. Geosyst.* **9**, Q01P23 (2008).
47. G. H. Haug, R. Tiedemann, R. Zahn, A. C. Ravelo, Role of Panama uplift on oceanic freshwater balance. *Geology* **29**, 207–210 (2001).
48. L. D. Keigwin Jr., Isotope paleoceanography of the Caribbean and east Pacific: Role of Panama uplift in late Neogene time. *Science* **217**, 350–353 (1982).
49. G. Keller, C. E. Zenker, S. M. Stone, Late neogene history of the Pacific-Caribbean gateway. *J. South Am. Earth Sci.* **2**, 73–108 (1989).
50. S. Steph, R. Tiedemann, M. Prange, J. Groeneveld, D. Nürnberg, L. Reuning, M. Schulz, G. H. Haug, Changes in Caribbean surface hydrography during the Pliocene shoaling of the Central American Seaway. *Paleoceanography* **21**, PA4221 (2006).
51. J. L. Teranes, D. H. Geary, B. E. Bemis, "The oxygen isotope record of seasonality in Neogene bivalves from the Central American Isthmus," in *Evolution and Environment in Tropical America*, J. B. C. Jackson, A. F. Budd, A. G. Coates, Eds. (University of Chicago Press, 1996), pp. 105–129.
52. A. M. Mestas-Núñez, P. Molnar, A mechanism for freshening the Caribbean Sea in pre-Ice Age time. *Paleoceanography* **29**, 508–517 (2014).
53. P. K. Weyl, The role of the oceans in climatic change: A theory of the ice ages. *Meteorol. Monogr.* **8**, 37–62 (1968).
54. A. M. Mestas-Núñez, D. B. Enfield, C. Zhang, Water vapor fluxes over the Intra-Americas Sea: Seasonal and interannual variability and associations with rainfall. *J. Climate* **20**, 1910–1922 (2007).
55. A. O'Dea, J. B. C. Jackson, H. Fortunato, J. T. Smith, L. D'Croz, K. G. Johnson, J. A. Todd, Environmental change preceded Caribbean extinction by 2 million years. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 5501–5506 (2007).
56. H. L. O. McClelland, P. D. Taylor, A. O'Dea, B. Okamura, Revising and refining the bryozoan *zs*-MART seasonality proxy. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **410**, 412–420 (2014).
57. R. Bonnefille, Cenozoic vegetation, climate changes and hominid evolution in tropical Africa. *Global Planet. Change* **72**, 390–411 (2010).
58. H. M. Liddy, S. J. Feakins, J. E. Tierney, Cooling and drying in northeast Africa across the Pliocene. *Earth Planet. Sci. Lett.* **449**, 430–438 (2016).
59. M. Brunet, A. Beauvilain, Y. Coppens, E. Heintz, A. H. E. Moutaye, D. Pilbeam, The first australopithecine 2,500 kilometres west of the Rift Valley (Chad). *Nature* **378**, 273–275 (1995).
60. A. Louchart, C. Mourer-Chauviré, H. T. Mackaye, A. Likius, P. Vignaud, M. Brunet, Les oiseaux du Pliocène inférieure du Jourab, Tchad, Afrique central. *Bull. Soc. Géol. Fr.* **175**, 413–421 (2004).
61. S. Leroy, L. Dupont, Development of vegetation and continental aridity in northwestern Africa during the Late Pliocene: The pollen record of ODP site 658. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **109**, 295–316 (1994).
62. L. Segalen, M. Renard, M. Pickford, B. Senut, I. Cojan, L. Le Callonnec, P. Rognon, Environmental and climatic evolution of the Namib Desert since the Middle Miocene: The contribution of carbon isotope ratios in ratite eggshells. *C. R. Geosci.* **334**, 917–924 (2002).
63. R. M. Forester, Pliocene-climate history of the western United States derived from lacustrine ostracodes. *Quat. Sci. Rev.* **10**, 133–146 (1991).
64. R. S. Thompson, Pliocene environments and climates in the western United States. *Quat. Sci. Rev.* **10**, 115–132 (1991).
65. M. Archer, S. J. Hand, H. Godthelp, "Tertiary environmental and biotic change in Australia," in *Paleoclimate and Evolution with Emphasis on Human Origins*, E. S. Vrba, G. H. Denton, T. C. Partridge, L. H. Burckle, Eds. (Yale Univ. Press, 1995), pp. 77–90.
66. J. M. Bowler, Aridity in Australia: Age, origin, and expression in aeolian landforms and sediments. *Earth Sci. Rev.* **12**, 279–310 (1976).
67. M. K. Macphail, Late Neogene climates in Australia: Fossil pollen- and spore-based estimates in retrospect and prospect. *Austral. J. Bot.* **45**, 425–464 (1997).
68. Z. An, J. E. Kutzbach, W. L. Prell, S. C. Porter, Evolution of Asian monsoons and phased uplift of the Himalaya-Tibet plateau since late Miocene times. *Nature* **411**, 62–66 (2001).
69. D. K. Rea, H. Snoeckx, L. H. Joseph, Late Cenozoic eolian deposition in the North Pacific: Asian drying, Tibetan uplift, and cooling of the northern hemisphere. *Paleoceanography* **13**, 215–224 (1998).
70. F. T. Kyte, M. Leinen, G. R. Heath, L. Zhou, Cenozoic sedimentation of the central North Pacific: Inferences from the elemental geochemistry of core LL44-GPC3. *Geochim. Cosmochim. Acta* **57**, 1719–1740 (1993).
71. A. Martínez-García, A. Rosell-Melé, S. L. Jaccard, W. Geibert, D. M. Sigman, G. H. Haug, Southern Ocean dust-climate coupling over the past four million years. *Nature* **476**, 312–315 (2011).
72. D. K. Rea, The paleoclimatic record provided by eolian deposition in the deep sea: The geologic history of wind. *Rev. Geophys.* **32**, 159–195 (1994).
73. S. A. Hovan, "Late Cenozoic atmospheric circulation intensity and climatic history recorded by eolian deposition in the eastern equatorial Pacific, Leg 138," in *Proc. Ocean Drill. Program Sci. Results 138*, N. G. Pisias, L. A. Mayer, T. R. Janacek, A. Palmer-Julson, T. H. van Andel, Eds., (Ocean Drilling Program, College Station, 1995), pp. 615–625.
74. J. K. Willenbring, D. J. Jerolmack, The null hypothesis: Globally steady rates of erosion, weathering fluxes and shelf sediment accumulation during Late Cenozoic mountain uplift and glaciation. *Terra Nova* **28**, 11–18 (2016).
75. F. Herman, J.-D. Champagnac, Plio-Pleistocene increase of erosion rates in mountain belts in response to climate change. *Terra Nova* **28**, 2–10 (2016).
76. F. Herman, D. Seward, P. G. Valla, A. Carter, B. Kohn, S. D. Willett, T. A. Ehlers, Worldwide acceleration of mountain erosion under a cooling climate. *Nature* **504**, 423–426 (2013).
77. W. W. Hay, J. L. Sloan II, C. N. Wold, Mass/age distribution and composition of sediments on the ocean floor and the global rate of sediment subduction. *J. Geophys. Res.* **93**, 14933–14940 (1988).
78. P. Molnar, Late Cenozoic increase in accumulation rates of terrestrial sediment: How might climate change have affected erosion rates? *Annu. Rev. Earth Planet. Sci.* **32**, 67–89 (2004).
79. P.-Z. Zhang, P. Molnar, W. R. Downs, Increased sedimentation rates and grain sizes 2–4 Myr ago due to the influence of climate change on erosion rates. *Nature* **410**, 891–897 (2001).
80. W. E. Galloway, T. L. Whiteaker, P. Ganey-Curry, History of Cenozoic North American drainage basin evolution, sediment yield, and accumulation in the Gulf of Mexico basin. *Geosphere* **7**, 938–973 (2011).
81. W. W. Hay, C. A. Shaw, C. N. Wold, Mass-balanced paleogeographic reconstructions. *Geol. Rundsch.* **78**, 207–242 (1989).
82. K. E. Murray, P. W. Reiners, S. N. Thomson, Rapid Pliocene–Pleistocene erosion of the central Colorado Plateau documented by apatite thermochronology from the Henry Mountains. *Geology* **44**, 483–486 (2016).

83. S. P. S. Gulick, J. M. Jaeger, A. C. Mix, H. Asahi, H. Bahlburg, C. L. Belanger, G. B. B. Berbel, L. Childress, E. Cowan, L. Drab, M. Forwick, A. Fukumural, S. Ge, S. Gupta, A. Kioka, S. Konno, L. J. LeVay, C. März, K. M. Matsuzaki, E. L. McClymont, C. Moy, J. Müller, A. Nakamura, T. Ojima, F. R. Ribeiro, K. D. Ridgway, O. E. Romero, A. L. Slagle, J. S. Stoner, G. St-Onge, I. Suto, M. D. Walczak, L. L. Worthington, I. Bailey, E. Enkelmann, R. Reece, J. M. Swartz, Mid-Pleistocene climate transition drives net mass loss from rapidly uplifting St. Elias Mountains, Alaska. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 15042–15047 (2015).
84. R. O. Lease, P. J. Haeussler, P. O'Sullivan, Changing exhumation patterns during Cenozoic growth and glaciation of the Alaska Range: Insights from detrital thermochronology and geochronology. *Tectonics* **35**, 934–955 (2016).
85. J. Kuhlemann, Post-collisional sediment budget of circum-Alpine basins (Central Europe). *Mem. Sci. Geol. Padova* **52**, 1–91 (2001).
86. J. Charreau, P.-H. Blard, N. Puchol, J.-P. Avouac, E. Lallier-Vergès, D. Bourlès, R. Braucher, A. Gallaud, R. Finkel, M. Jolivet, Y. Chen, P. Roy, Paleo-erosion rates in Central Asia since 9 Ma: A transient increase at the onset of Quaternary glaciations? *Earth Planet. Sci. Lett.* **304**, 85–92 (2011).
87. A. E. Blythe, D. W. Burbank, A. Carter, K. Schmidt, J. Putkonen, Plio-Quaternary exhumation history of the central Nepalese Himalaya: 1. Apatite and zircon fission track and apatite [U-Th]/He analyses. *Tectonics* **26**, TC3002 (2007).
88. K. W. Huntington, A. E. Blythe, K. V. Hodges, Climate change and Late Pliocene acceleration of erosion in the Himalaya. *Earth Planet. Sci. Lett.* **252**, 107–118 (2006).
89. M. J. Streule, A. Carter, M. P. Searle, J. M. Cottle, Constraints on brittle field exhumation of the Everest-Makalu section of the Greater Himalayan Sequence: Implications for models of crustal flow. *Tectonics* **31**, TC3010 (2012).

**Acknowledgments:** Comments by C. D. Bacon and C. Jaramillo helped this work. **Funding:** This work was supported by the Crafoord Foundation. **Competing interests:** The author declares that he has no competing interests. **Data and materials availability:** All data are available in the papers cited.

Submitted 21 September 2016  
Accepted 25 April 2017  
Published 14 June 2017  
10.1126/sciadv.1602320

**Citation:** P. Molnar, Comment (2) on "Formation of the Isthmus of Panama" by O'Dea *et al.* *Sci. Adv.* **3**, e1602320 (2017).

## Comment (2) on "Formation of the Isthmus of Panama" by O'Dea *et al.*

Peter Molnar

*Sci Adv* 3 (6), e1602320.  
DOI: 10.1126/sciadv.1602320

ARTICLE TOOLS	<a href="http://advances.sciencemag.org/content/3/6/e1602320">http://advances.sciencemag.org/content/3/6/e1602320</a>
REFERENCES	This article cites 82 articles, 26 of which you can access for free <a href="http://advances.sciencemag.org/content/3/6/e1602320#BIBL">http://advances.sciencemag.org/content/3/6/e1602320#BIBL</a>
PERMISSIONS	<a href="http://www.sciencemag.org/help/reprints-and-permissions">http://www.sciencemag.org/help/reprints-and-permissions</a>

Use of this article is subject to the [Terms of Service](#)

---

*Science Advances* (ISSN 2375-2548) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science Advances* is a registered trademark of AAAS.